



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Utility of phase angle to identify cachexia and assess mortality in end-stage renal disease

Seres, D., Mckeaveney, C., Adamson, G., Davenport, A., Farrington, K., Fouque, D., Kalanter-Zadeh, K., Mallett, J., Maxwell, P., Mullan, R., Nobel, H., O'Donoghue, D., Porter, S., Shields, J., Slee, A., Witham, M., & Reid, J. (2020). Utility of phase angle to identify cachexia and assess mortality in end-stage renal disease. *Current Developments in Nutrition*, 4(Supplement 2), Article 4141144. [https://doi.org/10.1093/cdn/nzaa055\\_029](https://doi.org/10.1093/cdn/nzaa055_029)

### Published in:

Current Developments in Nutrition

### Document Version:

Publisher's PDF, also known as Version of record

### Queen's University Belfast - Research Portal:

[Link to publication record in Queen's University Belfast Research Portal](#)

### Publisher rights

Copyright 2020 the authors.

This is an open access article published under a Creative Commons Attribution-NonCommercial-NoDerivs License

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

### General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

### Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

### Utility of Phase Angle to Identify Cachexia and Assess Mortality in End-Stage Renal Disease

David Seres,<sup>1</sup> Clare Mckeaveney,<sup>2</sup> Gary Adamson,<sup>3</sup> Andrew Davenport,<sup>4</sup> Ken Farrington,<sup>5</sup> Denis Fouque,<sup>6</sup> Kamyar Kalanter-Zadeh,<sup>7</sup> John Mallett,<sup>3</sup> Peter Maxwell,<sup>8</sup> Robert Mullan,<sup>9</sup> Helen Nobel,<sup>2</sup> Donal O'Donoghue,<sup>10</sup> Sam Porter,<sup>11</sup> Joanne Shields,<sup>12</sup> Adrian Slee,<sup>13</sup> Miles Witham,<sup>14</sup> and Joanne Reid<sup>15</sup>

<sup>1</sup>Columbia University Irving Medical Center; <sup>2</sup>Queen's University Belfast, UK; <sup>3</sup>Ulster University, Coleraine, UK; <sup>4</sup>University College London, United Kingdom; <sup>5</sup>University of Hertfordshire, Hatfield, UK; <sup>6</sup>University of Lyon, France; <sup>7</sup>University of California, Irvine; <sup>8</sup>Queens University Belfast & Regional Nephrology Unit, Belfast City Hospital, Belfast, UK; <sup>9</sup>Nephrology Department, Northern Health & Social Care Trust, UK; <sup>10</sup>Manchester University, UK; <sup>11</sup>University of Bournemouth; <sup>12</sup>Regional Nephrology Unit, Belfast City Hospital; <sup>13</sup>University College London; <sup>14</sup>NIHR Newcastle Biomedical Research Centre; and <sup>15</sup>Queen's University Belfast

**Objectives:** This cross-sectional analysis sought to identify cachexia and assess survival using phase angle (PA) in patients with end-stage renal disease (ESRD) receiving haemodialysis.

**Methods:** Patients receiving haemodialysis ( $n = 87$ , mean age 65.9  $\pm$  13.0) completed a Phase Angle (PA; 50 khz) measurement using bioelectrical impedance analysis. Cachexia variables were recorded according to Evans et al. definition (2008) including nutritional and functional measures (weight, Body Mass Index (BMI), Hand Grip Strength (HGS), Lean Tissue Mass (LTM), C-Reactive Protein

(CRP), serum albumin, haemoglobin, appetite (Functional Assessment of Anorexia/Cachexia Treatment (FAACT)) and fatigue (Functional Assessment of Chronic Illness Therapy (FACIT)). Survival was assessed at 12 months. Mann Whitney-U and Spearman correlation coefficient were conducted.

**Results:** The majority of patients completed follow up ( $n = 76$ ). Eleven patients had died. Mean PA was not statistically different between those identified as cachectic and non-cachectic according to Evans et al. (2008) definition or between those patients that survived and died. However, patients that survived had better mean scores of weight, BMI, HGS, CRP, serum albumin and fatigue (FACIT). In addition, LTM scores were significantly better in patients that survived ( $P < .01$ ). Appetite scores were also significantly better in patients that survived ( $P < .01$ ) and those without cachexia ( $P = .01$ ).

**Conclusions:** This study was part of a larger effort to clarify a phenotype of cachexia in ESRD. Unlike previous research, this study did not find PA useful in identifying patients at a higher risk of cachexia or death. However overall these patients had a very low mean PA. FAACT did discriminate between groups indicating self-reporting measurement tools of nutritional status were useful in identifying patients at a higher risk of cachexia and death. A larger sample and longer follow up is required to balance the limitations of this small study. Timing the administration of PA also requires consideration in future studies.

**Funding Sources:** Public Health Agency; Northern Ireland Kidney Research Fund.