



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Risks of digital screen time and recommendations for mitigating adverse outcomes in children and adolescents**

Pardhan, S., Parkin, J., Trott, M., & Driscoll, R. (2022). Risks of digital screen time and recommendations for mitigating adverse outcomes in children and adolescents. *Journal of School Health*, 92(8), 765-773. <https://doi.org/10.1111/josh.13170>

**Published in:**  
Journal of School Health

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

**Publisher rights**  
© 2022 American School Health Association.

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

**General rights**  
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

**Open Access**  
This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

# **Risks of digital screen time and recommendations for mitigating adverse outcomes in children and adolescents**

## **ABSTRACT**

### **BACKGROUND**

The COVID-19 pandemic caused an unprecedented move to emergency remote learning around the world, leading to increased digital screen time for children and adolescents. This review highlights the potential risk of increased screen time to the eye and general health and make recommendations to mitigate the risks posed.

### **METHODS**

A narrative review of evidence of increased digital time during the COVID-19 pandemic, the risks linked to increased screen time and offer possible steps to mitigate these in students.

### **RESULTS**

Digital screen time was found to have increased for children and adolescents in all the studies examined during the pandemic and data suggests that this has an impact on eye and general health. We discuss the associated risk factors and adverse outcomes associated with increased digital screen time.

### **CONCLUSIONS**

This review offers evidence of increased digital time, highlights some of the well-known and not so well-known risks linked to increased screen time and offer possible steps to mitigate these in children and adolescents during the pandemic, as well as offering schools and parents strategies to support the eye health of children and adolescents post-pandemic. We discuss a number of interventions to reduce the risk of eye strain, myopia, obesity and related diseases that have been shown to be linked to increased digital screen time.

**Keywords: Eye health, digital screen time, children, myopia**

The COVID-19 pandemic, originally reported in China, has spread rapidly around the world. In trying to reduce the spread of infection, governments around the world introduced emergency measures including restrictions on unnecessary movement and closure of schools. This has led to teaching courses being delivered online, with resulting increases in screen time and a reduction in time spent outdoors and physical activity.<sup>1</sup> According to the United Nations Educational, Scientific and Cultural Organization (UNESCO), approximately 1.37 billion students (80% of the world's student population) from more than 130 countries globally have been affected by these lockdown measures.<sup>2</sup> School closures caused an unprecedented move to emergency remote learning, defined as 'a temporary shift of instructional delivery to an alternate delivery mode due to crisis circumstances'<sup>3</sup>, with many countries adopting digital or e-learning approaches that have replaced face to face and classroom-based learning.

Whilst it is accepted that social isolation may be relieved by increased digital screen time, it is also very important to understand the potential risk to eye and general health of this increased screen time and ensure that a balance is achieved. Increased recreational screen time in schoolchildren (8-17 years old) has previously been linked to various adverse health conditions including eyesight

problems, an increase in obesity, insufficient sleep and inadequate physical activity<sup>4</sup>, and it is likely that local and national lockdowns due to the pandemic will exacerbate these issues.

## LITERATURE REVIEW

### Methods

The literature searches for this review were conducted in May 2021, and electronic databases (Scopus and PubMed) were searched using terms including “COVID”, “screen time” and “children”. Peer-reviewed articles which reported on screen time use during the COVID-19 pandemic were included, while irrelevant articles, those with no data on screen time and unavailable full texts were excluded.

### Evidence of increase in digital screen time globally

Evidence suggest that digital screen time has increased globally due to the COVID-19 pandemic. In Italy, digital screen time increased by  $4.85 \pm 2.40$  hours/day among children and adolescents (6-18 years old) during the COVID-19 lockdown.<sup>5</sup> Moore *et al.*<sup>6</sup> reported a mean screen time of 5.14 hours/day in younger children (5-11 years old) and 6.53 hours for older children (12-17 years old) with 88.7% of parents reporting that their children were exceeding the Canadian screen time guidelines of 2 hours per day. In Singapore, mean non-academic screen time increased from 1.61 hours to 3.15 hours/day during the national lockdown period in children aged 3-16 years old.<sup>7</sup> In Turkey, 71.7% of families surveyed reported an increase in children’s (6-13 years old) screen time ( $6.42 \pm 3.07$  hours/day during lockdown).<sup>8</sup> In Germany, a study found that recreational screen time among 4- to 17-year-olds increased from an average of 133.3 minutes to 194.5 minutes/day (an average increase of 61.2 minutes/day).<sup>9</sup> In Shanghai, children and adolescents (6-17 years) were found to have increased leisure screen time from an average of 170 to 450 minutes/week with a total screen time increasing from 610 to 2340 minutes per week.<sup>10</sup> In Tunisia, there was an increase of 111.11% in total screen time for children aged 5-12<sup>11</sup>, while in Saudi Arabia, mean digital device usage in children aged 10-18 increased from  $1.9 \pm 1.1$  hours to  $3.9 \pm 1.9$  hours/day, and the proportion using digital devices for more than 5 hours/day increased from 1.8% to 36.9%.<sup>12</sup> 44.6% of Chinese students (mean age 13) reported using digital devices for more than 5 hours/day for online study, which was associated with reports of irregular sleep patterns.<sup>13</sup> In the USA, recreational screen time among adolescents (14-18 years old) with autism spectrum disorder increased from 3.69 to 4.24 hours/day during the week, and from 5.84 to 7.39 hours/day at weekends.<sup>14</sup> A study in Chile found that toddlers and preschoolers (1-5 years old) increased their daily screen time from 1.66 to 3.05 hours/day during the pandemic.<sup>15</sup>

### Risks of increased digital screen time

**A. Digital eye strain.** Digital eye strain is characterized by eye discomfort during or after using a digital screen. The most common symptoms of digital eye strain are blurred vision, pain in the eyes, dry eyes and headaches. Even prior to COVID-19, studies report 59-95.8% of people experience at least one symptom of digital eye strain<sup>16, 17</sup>, with 56.5% of people reporting that their ocular symptoms have increased during the COVID-19 pandemic.<sup>17</sup> Here we highlight some of the reasons for digital eye strain.

**Uncorrected refractive error.** Whilst working on a computer does not make any refractive errors worse, any uncorrected errors that were previously tolerated may manifest themselves more readily. In some cases, it is a sign of normal ageing (affecting people around the age of 35-45 years)

when the lens inside the eye reduces its ability to focus for close distances (or loses its ability to 'accommodate'). It is a normal age-related change in the eye<sup>18-20</sup> and requires a correction for the near working distance.<sup>21-23</sup> In children, however this is not normally the case. However, if a child is strongly hyperopic (long-sighted) in one or both eyes, prolonged screen time may lead to eye strain due to the need for prolonged strain on the eye muscles as the child focusses for closer distance. This may be difficult to diagnose without an eye-test as long-sighted children are likely to have a good distance vision.

**Unstable binocular vision.** Binocular vision is being able to use both eyes adequately to create a single visual image. Uncorrected binocular vision problems increases the risk of blur and eye strain with screen time<sup>24-26</sup> Binocular vision problems, when the two eyes do not align properly to focus on one point in space, can create strain on the muscles in the eye as the individual is constantly trying to re-align the eyes to eliminate blurriness and double vision.<sup>27, 28</sup>

**Dry eyes.** Constant use of a computer screen can lead to significant discomfort due to the front part of the eyes drying out, especially if the eye 'forgets' to blink during periods of high concentration. Increased concentration during computer use can lead to a decrease in the number of blinks<sup>29, 30</sup> or to incomplete blinking<sup>31-33</sup>, both leading to reduced lubrication of the eyes resulting in dry eye symptoms.<sup>34</sup> The humidity of the room is also important as tear evaporation rate, eye comfort, tear stability, and production are adversely affected by dry air.<sup>35</sup>

**Screen distance.** The positioning of the computer screen is important, as advised by the Health and Safety Executive (HSE).<sup>36</sup> A desktop screen needs to be at a certain height and distance. A bigger desk area may be beneficial in order that the screen can be pushed further away if needed. It is possible that students learning at home during lockdown may not have been advised of the recommended distances especially if they are working in confined spaces.

**Hand-held devices.** Hand-held devices are near ubiquitous in US households, with 97% of children aged 0-8 having at least one smartphone and 75% having a tablet in their home.<sup>37</sup> Nearly half of 2- to 4-year-olds (46%) and 67% of 5- to 8-year-olds were reported to have their own device (tablet or smartphone) in 2020.<sup>37</sup> Most smartphones and tablets are viewed at shorter distances (between 20-40 cm). The screen and font size is also generally smaller compared to a desktop computer. The reduced working distance, smaller screen and font size will need more accommodative power in the eye with a higher risk of eye strain.

**Multiple devices.** It has been reported that children and adolescents often use several devices at once<sup>38, 39</sup>, for example to browse social media on their phone while watching content on another device.<sup>40</sup> A UK-based study of 816 adolescent females (mean age 12.8±0.8 years) reported that 59% of participants used more than one screen concurrently after school, with 65% using ≥2 screens concurrently in the evening, 36% before bed and 68% at weekends.<sup>41</sup> Switching between devices increases the strain on the eye by 22%, as this entails switching distances (and hence accommodative power of the eyes) between different devices.<sup>16</sup>

**B. Increased risk of myopia.** Increased digital screen time would lead to increase in near work and a reduction of outdoor activities, both of which are important risk factors for myopia.

**Increased near distance.** The exact role of increased near work, linked to digital screen time, as a risk factor for myopia onset and progression is still under debate. Some studies report an association between increased computer use and myopia<sup>42-44</sup>, whilst others fail to show that link to myopia prevalence.<sup>45</sup> A recent study reported a substantial myopic shift in children (6-13 years old) in China

as a result of home confinement.<sup>46</sup> Based on this, it is possible that earlier onset of myopia may occur due to increased digital screen time.<sup>47</sup>

**Decrease in outdoor activity.** It is undeniable that restrictive measures by governments have led to a decrease in outdoor activity. Increased outdoor activity is an important protective factor for the onset of myopia.<sup>48, 49</sup> Reports suggest that an additional 40 minutes per weekday of outdoor activity leads to a 23% reduction in the incidence of myopia.<sup>50</sup> Wu *et al.*<sup>51</sup> reported a 54% lower risk of myopia progression in children (6-7 years old) who spent more than 11 hours outdoors per week.

**C. Neck and shoulder strain.** It is recommended that the computer screen is at eye level and correct posture is vital.<sup>36</sup> Incorrect posture has been shown to lead to significant back problems.<sup>52</sup> The use of portable devices requires the need to incline our heads forward by an average of 27°, leading to an increased strain on the neck and shoulders.<sup>53-55</sup> A 27° incline forward of the head has been equated to an increased perceived weight of 18kg whilst a 60° angle looking at a mobile device equates to 27kg strain.<sup>25, 56</sup> In addition, texting using a mobile device has been found to be associated with a non-neutral posture of the cervical spine.<sup>57</sup>

**D. Increased sedentary time, overeating and obesity and associated higher risk of systemic diseases.** The restricted access to the outdoors and increased screen time during the pandemic, has typically led to increased sedentary time with reports suggesting a reduction of nearly 2.30 h/week of exercise time in children and adolescents.<sup>5</sup> A study from China found that children and adolescents (6-17 years) reduced their physical activity from an average of 540 to 105 minutes per week.<sup>10</sup> Conversely, a study on children in Germany found an increase of an average 36.2 minutes a day in total physical activity among children and young people (4-17 years), which the authors attributed to a substantial increase in habitual physical activities such as playing outside, gardening, housework, and walking and cycling.<sup>9</sup>

Increased screen time and sedentary behavior has been linked to obesity and increased body mass index (BMI) in youths<sup>58, 59</sup>, perhaps due to increased exposure to food advertising and passive food consumption.

Screen time has been found to be associated with increased dietary intake, reported to be due to distractions, interruption of physiologic food regulation, screen time as a conditioned cue to eat, disruption of memory formation, and the effects of the stress-induced food-reward system.<sup>60</sup> Using a smartphone during a meal can significantly affect the number of calories ingested, with one study finding an increase in caloric ingestion of 15% compared to eating without distracters.<sup>61</sup>

Poor eating habits have been linked to increased screen time<sup>62</sup>, more snack consumption and diet changes.<sup>63, 64</sup> Eating while viewing television has been shown to be positively associated with being overweight in children and adolescents (aged ≤ 18 years).<sup>65</sup> Regular snacking while watching television was associated with being overweight or obesity in pre-school children (4-6 years), as was screen time over 180 minutes per day.<sup>66</sup> In US adolescents (grades 9-12), excessive screen time whilst watching television, use of smartphones, tablets, computers and videogames was associated with a higher likelihood of obesity through increased sugary beverage consumption and reduction in physical activity.<sup>67</sup> In the Netherlands, children (10-14 years) who used screens for over 20 hours per week consumed more snacks than those who used them for 6 hours or less per week.<sup>64</sup>

It is also likely that the pandemic may have reduced the mental wellbeing of children and adults, leading to stress-eating of unhealthy foods leading to linked risk factors of weight gain. Unhealthy food habits increasing among obese children and adolescents (6-18 years old) have been reported in Italy during the pandemic.<sup>5</sup>

Increased BMI in childhood has been demonstrated to increase the likelihood of obesity in adulthood<sup>68</sup>, which in turn increases the risk of developing non-communicable diseases. Sedentary behavior has been positively associated with increased risk of disease later in life<sup>69</sup>, including Type 2 diabetes<sup>70</sup>, diabetic retinopathy<sup>71</sup>, cancer<sup>72-74</sup>, dementia<sup>75</sup>, and cardiovascular disease<sup>70</sup>. Low levels of physical activity, low cardiorespiratory fitness and sedentary behavior are significantly associated with the development of metabolic syndrome.<sup>76</sup>

**E. Disturbed sleep patterns.** Screen time has been linked to poor sleep in children and adolescents. A systematic review (1999-2014) reported that screen time was associated with poorer sleep outcomes.<sup>77</sup> Another systematic review of studies in children and adolescents aged 6-19 years reported significant associations between the presence of a media device in the sleep environment near bedtime and inadequate sleep quality, poor sleep quality, and excessive daytime sleepiness.<sup>78</sup> A Spanish study found that 44% of children aged between 1 and 14 years had over 120 minutes per day of leisure screen time, which was associated with short sleep duration.<sup>79</sup> A study of American teenagers found that screen device usage of 5 or more hours per day resulted in 80% higher odds of inadequate sleep compared to those who did not use devices (OR = 1.79, 95% CI: 1.54, 2.08).<sup>67</sup> Smartphone use in bed was found to be a strong predictor for sleep problems including prolonged sleep latency and short sleep duration in adolescents.<sup>80</sup> In children under 5, higher levels of daily screen time were associated with shorter sleep duration and more night awakenings.<sup>81</sup>

**F. Mental Health.** While moderate engagement in digital activities has found not to be harmful<sup>82</sup>, and it is clearly important to appreciate that using screens to maintain relationships with friends would mitigate feelings of isolation<sup>83, 84</sup>, research also suggest that high screen exposure increases depression among youths.<sup>59, 85</sup> Poorer mental health has also been shown among children and adolescents using screens for more than 2-3 hours per day<sup>86</sup>, with severe depressive symptoms being positively associated with higher levels of screen time.<sup>87</sup> Research has also linked problematic smart phone use (i.e. use associated with at least some element of dysfunctional use; examples include anxiety when a phone is not available, or neglect of other activities<sup>88, 89</sup>) to self-reported anxiety, insomnia, increased perceived stress, poor educational attainment<sup>90</sup> and decreased overall quality of life.<sup>91</sup> Excessive screen time has also been found to exacerbate ADHD-related behavior.<sup>92</sup> In children under 5 years of age, touch screen device use has been found to be more damaging than beneficial, and has been linked to emotional problems and attention problems.<sup>93</sup>

Primary caregivers of Portuguese children aged 6 to 82 months reported that screen exposure time increased during COVID-19 confinement, and this was positively correlated with behavioral and emotional problems, with attention problems and parent challenges affected most by increased screen time exposure.<sup>94</sup>

**G. Online gaming and other activity.** In addition, digital screen exposure has also increased for online gaming and streaming activity during the COVID-19 lockdown in different countries.<sup>95-100</sup>

### Long-term behavior change

Although school closures may be short-lived, it is possible that ease of online access to schoolwork may increase the widespread acceptance of online classes and increased use of digital screen time in the longer term, leading to behavioral changes that persist after the pandemic.

## IMPLICATIONS FOR SCHOOL HEALTH

### Mitigation

**Better ergonomics.** It is important that the computer is set up correctly in order to reduce eye and back strain. The screen should be at arms length away with the eyes centered in the top third of the screen. The head should be looking straight ahead. The seat adjusted so that feet are flat on the ground, neck and back straight, forearms same height as the desk. It is important to ensure that the screen is big enough for all the data and that access to the mouse and keyboard is comfortable. The screen should be positioned to avoid glare and reflection and the brightness and contrast of the screen adjusted to ensure they are comfortable. Following correct ergonomic advice and performing eye exercises for 4 weeks was demonstrated to reduce eye fatigue in children (aged 6 to 15 years) attending online classes during the COVID-19 pandemic.<sup>101</sup>

**Adequate visual function, blinking and good eye habits.** It is vital to develop good eye habits when using the screen, both during the pandemic and also beyond it. It is important that any uncorrected vision problems are adequately dealt with by an optometrist, ophthalmologist, or eyecare provider. Correct spectacle prescription for the screen distance and binocular vision anomalies should be addressed by exercises, spectacles, and other optical corrections. Adequate correction of binocular vision anomalies has been shown to reduce ocular symptoms of digital eye strain.<sup>102, 103</sup> Blinking exercises for people who suffer from dry eyes and who forget to blink are very useful.<sup>104</sup> A reminder on the computer every hour to forcibly blink ten times every hour can help. For those with more severe dry eyes, artificial tears and omega 3 food additives may be helpful. Trials have shown that interventions aimed at modifying diet and increasing physical activity can improve dry eye disease status<sup>105</sup>, with omega 3 supplements resulting in reduced tear osmolarity and increased tear stability in adults with dry eye disease.<sup>106, 107</sup>

The HSE advises short breaks often rather than longer ones less often. Five to ten minutes every hour is better than 20 minutes every 2 hours.<sup>36, 53</sup> A 20-20-20 rule is also useful- a 20 second break every 20 minutes to gaze a distance at least 20 feet away. Frequent short breaks can relax accommodative and binocular vision and encourage normal blinking therefore improving the tear quality. Software which instructs users to blink at regular intervals while using computers has been demonstrated to increase blink rate and improve dry eye symptoms in adults.<sup>108</sup>

**Smart use of digital screen time by users and schools.** Ensure time spent on digital devices are maximized for learning and less digital time is used for recreational activities. It would be important for government agencies to continue to engage with schools to shape a holistic home-based learning curriculum that encourages creative learning not just from reading and study at home, but also include frequent breaks and indoor physical or household activities such as cooking, baking, and cleaning.

**Limiting screen time.** A daily schedule setting time for when devices can be used would be beneficial. The World Health Organization's guidelines on physical activity, sedentary behavior, and sleep recommends less than 1 hour of sedentary screen time for children 1-5 years of age.<sup>109</sup> The continuous use of digital devices for non-educational purposes should be limited to <15 minutes per day and a cumulative duration of <1 hour a day. The American Academy of Pediatrics recommends restricting screen time to 1 hour per day of high-quality content for children 18 months to 5 years of age and suggests consistent limits for those 6 years of age and above should have 8-10 hours of day screen-free time.<sup>110</sup>

**Increased public awareness.** Public education to increase parent awareness about the effects of increased screen time is important over the long term.

**Parents and Teachers as role models.** Parents can act as role models. It is known that screen time exposure is related to family behaviors. An increase in physically active behavior such as playing indoors (if possible), baking, cooking, craft activities would benefit.<sup>111-114</sup> Xu *et al.*<sup>115</sup> found family roles to play an important role in promoting physical activity. Encouragement and support from parents can increase time spent by their children undergoing physical activity, and reducing parents screen time can lead to a decrease in their children's screen time.<sup>115</sup> Parents need to understand the importance of maintaining good eye habits during the pandemic lockdown and beyond, including frequent breaks from near work and limiting recreational screen time. Parents can reduce the detrimental effects of digital screen time by monitoring recreational usage, setting reminders to disconnect and ensuring that time spent on digital devices are maximized for learning.

**Increased time outdoors.** Working within government rules during the pandemic, allowable time outdoors should be fully utilized. Increased physical activity and outdoor play has been shown to lead to longer sleep in preschoolers<sup>81, 116</sup>, reduced obesity and also a reduced risk of myopia. Outdoor activity time has also been linked to pediatric dry eye disease, with a study of Korean school children (7 to 12 years old) finding that increased outdoor activity time reduced the rate of dry eye disease.<sup>117</sup>

## Conclusions

In this review we report increased digital time usage in children and adolescents during the COVID-19 pandemic, the associated risk factors and adverse outcomes and provide a number of interventions to reduce the risk of eye strain, myopia, obesity and related diseases that have been shown to be linked to increased digital screen time. Through increasing awareness of the risks associated with high levels of digital screen use and sharing strategies to reduce the negative effects, teachers and parents will be encouraged to enhance the health and wellbeing of children and adolescents in the pandemic and beyond.

## Human Subjects Approval Statement

Preparation of this paper did not involve primary research or data collection involving human subjects, and therefore, no institutional review board examination or approval was required.

## References



## REFERENCES

1. Stockwell S, Trott M, Tully M, et al. Changes in physical activity and sedentary behaviours from before to during the COVID-19 pandemic lockdown: A systematic review. *BMJ Open Sport Exerc Med.* 2021;7(1). doi: 10.1136/bmjsem-2020-000960.
2. UNESCO. 1.37 billion students now home as COVID-19 school closures expand, ministers scale up multimedia approaches to ensure learning continuity. 2021. Available from: <https://en.unesco.org/news/137-billion-students-now-home-covid-19-school-closures-expand-ministers-scale-multimedia>. Accessed 01/09/2021.
3. Hodges C, Moore S, Lockee B, Trust T, Bond M. The Difference Between Emergency Remote Teaching and Online Learning. 2020.
4. Tambalis KD, Panagiotakos DB, Psarra G, Sidossis LS. Screen time and its effect on dietary habits and lifestyle among schoolchildren. *Cent Eur J Public Health.* 2020;28(4):260-266. doi: 10.21101/cejph.a6097.
5. Pietrobelli A, Pecoraro L, Ferruzzi A, et al. Effects of COVID-19 Lockdown on Lifestyle Behaviors in Children with Obesity Living in Verona, Italy: A Longitudinal Study. *Obesity.* 2020;28(8):1382-1385. doi: 10.1002/oby.22861.
6. Moore SA, Faulkner G, Rhodes RE, et al. Impact of the COVID-19 virus outbreak on movement and play behaviours of Canadian children and youth: A national survey. *Int J Behav Nutr Phys Act.* 2020;17(1). doi: 10.1186/s12966-020-00987-8.
7. Lim MTC, Ramamurthy MB, Aishworiya R, et al. School closure during the coronavirus disease 2019 (COVID-19) pandemic – Impact on children's sleep. *Sleep Med.* 2021;78:108-114. doi: 10.1016/j.sleep.2020.12.025.
8. Ozturk Eyimaya A, Yalçın Irmak A. Relationship between parenting practices and children's screen time during the COVID-19 Pandemic in Turkey. *J Pediatr Nurs.* 2021;56:24-29. doi: 10.1016/j.pedn.2020.10.002.

9. Schmidt SCE, Anedda B, Burchartz A, et al. Physical activity and screen time of children and adolescents before and during the COVID-19 lockdown in Germany: a natural experiment. *Sci Rep.* 2020;10(1). doi: 10.1038/s41598-020-78438-4.
10. Xiang M, Zhang Z, Kuwahara K. Impact of COVID-19 pandemic on children and adolescents' lifestyle behavior larger than expected. *Prog Cardiovasc Dis.* 2020;63(4):531-532. doi: 10.1016/j.pcad.2020.04.013.
11. Abid R, Ammar A, Maaloul R, Souissi N, Hammouda O. Effect of COVID-19-related home confinement on sleep quality, screen time and physical activity in tunisian boys and girls: A survey. *Int J Environ Res Public Health.* 2021;18(6):1-12. doi: 10.3390/ijerph18063065.
12. Mohan A, Sen P, Shah C, Jain E, Jain S. Prevalence and risk factor assessment of digital eye strain among children using online e-learning during the COVID-19 pandemic: Digital eye strain among kids (DESK study-1). *Indian J Ophthalmol.* 2021;69(1):140-144. doi: 10.4103/ijo.IJO\_2535\_20.
13. Guo Y-, Liao M-, Cai W-, et al. Physical activity, screen exposure and sleep among students during the pandemic of COVID-19. *Sci Rep.* 2021;11(1). doi: 10.1038/s41598-021-88071-4.
14. Garcia JM, Lawrence S, Brazendale K, Leahy N, Fukuda D. Brief report: The impact of the COVID-19 pandemic on health behaviors in adolescents with Autism Spectrum Disorder. *Disability and Health Journal.* 2021;14(2):101021. doi: <https://doi.org/10.1016/j.dhjo.2020.101021>.
15. Aguilar-Farias N, Toledo-Vargas M, Miranda-Marquez S, et al. Sociodemographic predictors of changes in physical activity, screen time, and sleep among toddlers and preschoolers in chile during the covid-19 pandemic. *Int J Environ Res Public Health.* 2021;18(1):1-13. doi: 10.3390/ijerph18010176.
16. The Vision Council. Digital Eye Strain Report "Eyes Overexposed: The Digital Device Dilemma". 2016.
17. Bahkir FA, Grandee SS. Impact of the COVID-19 lockdown on digital device-related ocular health. *Indian J Ophthalmol.* 2020;68(11):2378-2383. doi: 10.4103/ijo.IJO\_2306\_20.

18. Martin H, Guthoff R, Terwee T, Schmitz K-. Comparison of the accommodation theories of Coleman and of Helmholtz by finite element simulations. *Vis Res.* 2005;45(22):2910-2915. doi: 10.1016/j.visres.2005.05.030.
19. Glasser A, Kaufman PL. The mechanism of accommodation in primates. *Ophthalmology.* 1999;106(5):863-872. doi: 10.1016/S0161-6420(99)00502-3.
20. Charman WN. The eye in focus: Accommodation and presbyopia. *Clin Exp Optom.* 2008;91(3):207-225. doi: 10.1111/j.1444-0938.2008.00256.x.
21. Collier JD, Rosenfield M. Accommodation and convergence during sustained computer work. *Optometry.* 2011;82(7):434-440. doi: 10.1016/j.optm.2010.10.013.
22. Hashemi H, Saatchi M, Yekta A, et al. High prevalence of asthenopia among a population of university students. *J Ophthalmic Vis Res.* 2019;14(4):474-482. doi: 10.18502/jovr.v14i4.5455.
23. Sánchez-Brau M, Domenech-Amigot B, Brocal-Fernández F, Quesada-Rico JA, Seguí-Crespo M. Prevalence of Computer Vision Syndrome and Its Relationship with Ergonomic and Individual Factors in Presbyopic VDT Workers Using Progressive Addition Lenses. *Int J Environ Res Public Health.* 2020;17(3). doi: 10.3390/ijerph17031003.
24. Mocci F, Serra A, Corrias GA. Psychological factors and visual fatigue in working with video display terminals. *Occup Environ Med.* 2001;58(4):267-271. doi: 10.1136/oem.58.4.267.
25. Ang C, Dinevski D, Vlasak N, Kok A. Lens Study: Taking the Strain. *Optician.* 2017;253.
26. Wick B, Morse S. Accommodative accuracy to video display monitors. *Optometry Vision Sci.* 2002;79(12). doi: 10.1097/00006324-200212001-00413.
27. Jaiswal S, Asper L, Long J, Lee A, Harrison K, Golebiowski B. Ocular and visual discomfort associated with smartphones, tablets and computers: what we do and do not know. *Clin Exp Optom.* 2019;102(5):463-477. doi: 10.1111/cxo.12851.
28. Yammouni R, Evans BJW. Is reading rate in digital eyestrain influenced by binocular and accommodative anomalies?. *J Optom.* 2020. doi: 10.1016/j.optom.2020.08.006.

29. Rosenfield M, Jahan S, Nunez K, Chan K. Cognitive demand, digital screens and blink rate. *Comput Hum Behav.* 2015;51(PA):403-406. doi: 10.1016/j.chb.2015.04.073.
30. Patel P, Henderson R, Bradley L, Galloway B, Hunter L. Effect of visual display unit use on blink rate and tear stability. *Optometry Vision Sci.* 1991;68(11):888-892. doi: 10.1097/00006324-199111000-00010.
31. Argilés M, Cardona G, Pérez-Cabré E, Rodríguez M. Blink rate and incomplete blinks in six different controlled hard-copy and electronic reading conditions. *Invest Ophthalmol Vis Sci.* 2015;56(11):6679-6685. doi: 10.1167/iovs.15-16967.
32. McMonnies CW. Incomplete blinking: Exposure keratopathy, lid wiper epitheliopathy, dry eye, refractive surgery, and dry contact lenses. *Contact Lens Anterior Eye.* 2007;30(1):37-51. doi: 10.1016/j.clae.2006.12.002.
33. Hirota M, Uozato H, Kawamorita T, Shibata Y, Yamamoto S. Effect of incomplete blinking on tear film stability. *Optom Vis Sci.* 2013;90(7):650-657. doi: 10.1097/OPX.0b013e31829962ec.
34. Golebiowski B, Long J, Harrison K, Lee A, Chidi-Egboka N, Asper L. Smartphone Use and Effects on Tear Film, Blinking and Binocular Vision. *Curr Eye Res.* 2020;45(4):428-434. doi: 10.1080/02713683.2019.1663542.
35. Abusharha AA, Pearce EI. The effect of low humidity on the human tear film. *Cornea.* 2013;32(4):429-434. doi: 10.1097/ICO.0b013e31826671ab.
36. Health and Safety Executive. Health and Safety (Display Screen Equipment) Regulations. 1992.
37. Rideout V, Robb MB. The common sense census. Media use by kids age zero to eight. 2020.
38. Jago R, Sebire SJ, Gorely T, Cillero IH, Biddle SJH. "I'm on it 24/7 at the moment": A qualitative examination of multi-screen viewing behaviours among UK 10-11 year olds. *Int J Behav Nutr Phys Act.* 2011;8. doi: 10.1186/1479-5868-8-85.

39. Toh SH, Howie EK, Coenen P, Straker LM. "From the moment i wake up i will use it...every day, very hour": A qualitative study on the patterns of adolescents' mobile touch screen device use from adolescent and parent perspectives. *BMC Pediatr.* 2019;19(1). doi: 10.1186/s12887-019-1399-5.
40. Ofcom. Children and parents: media use and attitudes report. 2021.
41. Harrington DM, Ioannidou E, Davies MJ, et al. Concurrent screen use and cross-sectional association with lifestyle behaviours and psychosocial health in adolescent females. *Acta Paediatr.* 2021;110(7):2164-2170. doi: <https://doi.org/10.1111/apa.15806>.
42. Enthoven CA, Tideman JWL, Polling JR, Yang-Huang J, Raat H, Klaver CCW. The impact of computer use on myopia development in childhood: The Generation R study. *Prev Med.* 2020;132:105988. doi: 10.1016/j.ypmed.2020.105988.
43. McCrann S, Loughman J, Butler JS, Paudel N, Flitcroft DI. Smartphone use as a possible risk factor for myopia. *Clin Exp Optom.* 2020. doi: 10.1111/cxo.13092.
44. Alvarez-Peregrina C, Sánchez-Tena MÁ, Martínez-Perez C, Villa-Collar C. The Relationship Between Screen and Outdoor Time With Rates of Myopia in Spanish Children. *Front Public Health.* 2020;8. doi: 10.3389/fpubh.2020.560378.
45. Lanca C, Saw S-. The association between digital screen time and myopia: A systematic review. *Ophthalmic Physiol Opt.* 2020;40(2):216-229. doi: 10.1111/opo.12657.
46. Wang J, Li Y, Musch DC, et al. Progression of Myopia in School-Aged Children after COVID-19 Home Confinement. *JAMA Ophthalmol.* 2021. doi: 10.1001/jamaophthalmol.2020.6239.
47. Klaver CCW, Polling JR, Enthoven CA. 2020 as the Year of Quarantine Myopia. *JAMA Ophthalmol.* 2021. doi: 10.1001/jamaophthalmol.2020.6231.
48. Xiong S, Sankaridurg P, Naduvilath T, et al. Time spent in outdoor activities in relation to myopia prevention and control: a meta-analysis and systematic review. *Acta Ophthalmol.* 2017;95(6):551-566. doi: 10.1111/aos.13403.

49. Eppenberger LS, Sturm V. The role of time exposed to outdoor light for myopia prevalence and progression: A literature review. *Clin Ophthalmol*. 2020;14:1875-1890. doi: 10.2147/OPTH.S245192.
50. He M, Xiang F, Zeng Y, et al. Effect of time spent outdoors at school on the development of myopia among children in China a randomized clinical trial. *JAMA*. 2015;314(11):1142-1148. doi: 10.1001/jama.2015.10803.
51. Wu P-, Chen C-, Lin K-, et al. Myopia Prevention and Outdoor Light Intensity in a School-Based Cluster Randomized Trial. *Ophthalmology*. 2018;125(8):1239-1250. doi: 10.1016/j.ophtha.2017.12.011.
52. NIOSH. Musculoskeletal disorders and workplace factors: a critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and low back. *Department of Health and Human Services, DHHS*. 1997;97-141.
53. Sheppard AL, Wolffsohn JS. Digital eye strain: Prevalence, measurement and amelioration. *BMJ Open Ophthalmology*. 2018;3(1). doi: 10.1136/bmjophth-2018-000146.
54. American Optometric Association. Computer Vision Syndrome. 2017.
55. Rosenfield M. Computer Vision Syndrome (aka digital eye strain). 2016;17.
56. Hansraj KK. Assessment of stresses in the cervical spine caused by posture and position of the head. *Surg Technol Int*. 2014;25:277-279.
57. Xie YF, Szeto G, Madeleine P, Tsang S. Spinal kinematics during smartphone texting – A comparison between young adults with and without chronic neck-shoulder pain. *Appl Ergon*. 2018;68:160-168. doi: 10.1016/j.apergo.2017.10.018.
58. Elgar FJ, Roberts C, Moore L, Tudor-Smith C. Sedentary behaviour, physical activity and weight problems in adolescents in Wales. *Public Health*. 2005;119(6):518-524. doi: 10.1016/j.puhe.2004.10.011.
59. Stiglic N, Viner RM. Effects of screentime on the health and well-being of children and adolescents: a systematic review of reviews. *BMJ Open*. 2019;9(1):e023191. doi: 10.1136/bmjopen-2018-023191.

60. Marsh S, Ni Mhurchu C, Maddison R. The non-advertising effects of screen-based sedentary activities on acute eating behaviours in children, adolescents, and young adults. A systematic review. *Appetite*. 2013;71:259-273. doi: 10.1016/j.appet.2013.08.017.
61. Gonçalves, Renata Fiche da Mata, Barreto Dda, Monteiro PI, et al. Smartphone use while eating increases caloric ingestion. *Physiol Behav*. 2019;204:93-99. doi: 10.1016/j.physbeh.2019.02.021.
62. Mattioli AV, Ballerini Puviani M, Nasi M, Farinetti A. COVID-19 pandemic: the effects of quarantine on cardiovascular risk. *Eur J Clin Nutr*. 2020;74(6):852-855. doi: 10.1038/s41430-020-0646-z.
63. Tsujiguchi H, Hori D, Kambayashi Y, et al. Relationship between screen time and nutrient intake in Japanese children and adolescents: a cross-sectional observational study. *Environmental health and preventive medicine*. 2018;23(1):34-42. doi: 10.1186/s12199-018-0725-0.
64. Berentzen NE, Smit HA, van Rossem L, et al. Screen time, adiposity and cardiometabolic markers: mediation by physical activity, not snacking, among 11-year-old children. *International Journal of Obesity*. 2014;38(10):1317-1323. doi: 10.1038/ijo.2014.110.
65. Ghobadi S, Hassanzadeh-Rostami Z, Salehi-Marzijarani M, et al. Association of eating while television viewing and overweight/obesity among children and adolescents: a systematic review and meta-analysis of observational studies. *Obes Rev*. 2018;19(3):313-320. doi: 10.1111/obr.12637.
66. Kurspahić-Mujčić A, Mujčić A. Factors associated with overweight and obesity in preschool children. *Med Glas*. 2020;17(2):538-543. doi: 10.17392/1175-20.
67. Kenney EL, Gortmaker SL. United States Adolescents' Television, Computer, Videogame, Smartphone, and Tablet Use: Associations with Sugary Drinks, Sleep, Physical Activity, and Obesity. *J Pediatr*. 2017;182:144-149. doi: 10.1016/j.jpeds.2016.11.015.
68. Rundle AG, Factor-Litvak P, Suglia SF, et al. Tracking of Obesity in Childhood into Adulthood: Effects on Body Mass Index and Fat Mass Index at Age 50. *Childhood Obesity*. 2020;16(3):226-233. doi: 10.1089/chi.2019.0185.

69. Patterson R, McNamara E, Tainio M, et al. Sedentary behaviour and risk of all-cause, cardiovascular and cancer mortality, and incident type 2 diabetes: a systematic review and dose response meta-analysis. *Eur J Epidemiol*. 2018;33(9):811-829. doi: 10.1007/s10654-018-0380-1.
70. Bailey DP, Hewson DJ, Champion RB, Sayegh SM. Sitting Time and Risk of Cardiovascular Disease and Diabetes: A Systematic Review and Meta-Analysis. *Am J Prev Med*. 2019;57(3):408-416. doi: 10.1016/j.amepre.2019.04.015.
71. Ren C, Liu W, Li J, Cao Y, Xu J, Lu P. Physical activity and risk of diabetic retinopathy: a systematic review and meta-analysis. *Acta Diabetol*. 2019;56(8):823-837. doi: 10.1007/s00592-019-01319-4.
72. Biller VS, Leitzmann MF, Sedlmeier AM, Berger FF, Ortmann O, Jochem C. Sedentary behaviour in relation to ovarian cancer risk: a systematic review and meta-analysis. *Eur J Epidemiol*. 2021. doi: 10.1007/s10654-020-00712-6.
73. Chan DSM, Abar L, Cariolou M, et al. World Cancer Research Fund International: Continuous Update Project—systematic literature review and meta-analysis of observational cohort studies on physical activity, sedentary behavior, adiposity, and weight change and breast cancer risk. *Cancer Causes Control*. 2019;30(11):1183-1200. doi: 10.1007/s10552-019-01223-w.
74. Berger FF, Leitzmann MF, Hillreiner A, et al. Sedentary behavior and prostate cancer: A systematic review and meta-analysis of prospective cohort studies. *Cancer Prev Res*. 2019;12(10):675-687. doi: 10.1158/1940-6207.CAPR-19-0271.
75. Yan S, Fu W, Wang C, et al. Association between sedentary behavior and the risk of dementia: a systematic review and meta-analysis. *Transl Psychiatry*. 2020;10(1). doi: 10.1038/s41398-020-0799-5.
76. De Oliveira RG, Guedes DP. Physical activity, sedentary behavior, cardiorespiratory fitness and metabolic syndrome in adolescents: Systematic review and meta-analysis of observational evidence. *PLoS ONE*. 2016;11(12). doi: 10.1371/journal.pone.0168503.
77. Hale L, Guan S. Screen time and sleep among school-aged children and adolescents: A systematic literature review. *Sleep Medicine Reviews*. 2015;21:50-58. doi: 10.1016/j.smr.2014.07.007.



78. Carter B, Rees P, Hale L, Bhattacharjee D, Paradkar MS. Association Between Portable Screen-Based Media Device Access or Use and Sleep Outcomes: A Systematic Review and Meta-analysis. *JAMA pediatrics*. 2016;170(12):1202-1208. doi: 10.1001/jamapediatrics.2016.2341.
79. Cartanyà-Hueso À, Lidón-Moyano C, Martín-Sánchez JC, et al. Association of screen time and sleep duration among Spanish 1-14 years old children. *Paediatr Perinat Epidemiol*. 2021;35(1):120-129. doi: 10.1111/ppe.12695.
80. Kater M-, Schlarb AA. Smartphone usage in adolescents – motives and link to sleep disturbances, stress and sleep reactivity. *Somnologie*. 2020;24(4):245-252. doi: 10.1007/s11818-020-00272-7.
81. Janssen X, Martin A, Hughes AR, Hill CM, Kotronoulas G, Hesketh KR. Associations of screen time, sedentary time and physical activity with sleep in under 5s: A systematic review and meta-analysis. *Sleep Med Rev*. 2020;49. doi: 10.1016/j.smr.2019.101226.
82. Przybylski AK, Weinstein N. A Large-Scale Test of the Goldilocks Hypothesis: Quantifying the Relations Between Digital-Screen Use and the Mental Well-Being of Adolescents. *Psychol Sci*. 2017;28(2):204-215. doi: 10.1177/0956797616678438.
83. Laursen B, Bukowski WM, Aunola K, Nurmi J-. Friendship moderates prospective associations between social isolation and adjustment problems in young children. *Child Dev*. 2007;78(4):1395-1404. doi: 10.1111/j.1467-8624.2007.01072.x.
84. Williams KD, Cheung CK, Choi W. Cyberostracism: effects of being ignored over the Internet. *J Pers Soc Psychol*. 2000;79(5):748-762. doi: 10.1037//0022-3514.79.5.748 [doi].
85. Kremer P, Elshaug C, Leslie E, Toumbourou JW, Patton GC, Williams J. Physical activity, leisure-time screen use and depression among children and young adolescents. *J Sci Med Sport*. 2014;17(2):183-187. doi: 10.1016/j.jsams.2013.03.012.
86. Hoare E, Milton K, Foster C, Allender S. The associations between sedentary behaviour and mental health among adolescents: A systematic review. *Int J Behav Nutr Phys Act*. 2016;13(1). doi: 10.1186/s12966-016-0432-4.

87. Mougharbel F, Goldfield GS. Psychological Correlates of Sedentary Screen Time Behaviour Among Children and Adolescents: a Narrative Review. *Curr Obesity Rep.* 2020;9(4):493-511. doi: 10.1007/s13679-020-00401-1.
88. Lin Y-, Chiang C-, Lin P-, et al. Proposed diagnostic criteria for Smartphone addiction. *PLoS ONE.* 2016;11(11). doi: 10.1371/journal.pone.0163010.
89. Bianchi A, Phillips JG. Psychological predictors of problem mobile phone use. *Cyberpsychol Behav.* 2005;8(1):39-51. doi: 10.1089/cpb.2005.8.39.
90. Sohn SY, Rees P, Wildridge B, Kalk NJ, Carter B. Prevalence of problematic smartphone usage and associated mental health outcomes amongst children and young people: a systematic review, meta-analysis and GRADE of the evidence. *BMC Psychiatry.* 2019;19(1):356. doi: 10.1186/s12888-019-2350-x.
91. Fischer-Grote L, Kothgassner OD, Felnhofer A. The impact of problematic smartphone use on children's and adolescents' quality of life: A systematic review. *Acta Paediatr Int J Paediatr.* 2021;110(5):1417-1424. doi: 10.1111/apa.15714.
92. Lissak G. Adverse physiological and psychological effects of screen time on children and adolescents: Literature review and case study. *Environ Res.* 2018;164:149-157. doi: 10.1016/j.envres.2018.01.015.
93. Lin H, Chen K, Chou W, et al. Prolonged touch screen device usage is associated with emotional and behavioral problems, but not language delay, in toddlers. *Infant Behavior and Development.* 2020;58:101424. doi: 10.1016/j.infbeh.2020.101424.
94. Monteiro R, Rocha NB, Fernandes S. Are Emotional and Behavioral Problems of Infants and Children Aged Younger Than 7 Years Related to Screen Time Exposure During the Coronavirus Disease 2019 Confinement? An Exploratory Study in Portugal. *Front Psychol.* 2021;12. doi: 10.3389/fpsyg.2021.590279.
95. King DL, Delfabbro PH, Billieux J, Potenza MN. Problematic online gaming and the COVID-19 pandemic. *J Behav Addict.* 2020;9(2):184-186. doi: 10.1556/2006.2020.00016.

96. Oflu A, Bükülmez A, Elmas E, Tahta EG, Çeleğen M. Comparison of screen time and digital gaming habits of turkish children before and during the coronavirus disease 2019 pandemic. *Turk Pediatr Ars.* 2021;56(1):22-26. doi: 10.14744/TurkPediatriArs.2020.41017.
97. Paschke K, Austermann MI, Simon-Kutscher K, Thomasius R. Adolescent gaming and social media usage before and during the COVID-19 pandemic. *Sucht.* 2021;67(1):13-22. doi: 10.1024/0939-5911/a000694.
98. Teng Z, Pontes HM, Nie Q, Griffiths MD, Guo C. Depression and anxiety symptoms associated with internet gaming disorder before and during the COVID-19 pandemic: A longitudinal study. *J Behav Addict.* 2021;10(1):169-180. doi: 10.1556/2006.2021.00016.
99. Zhu S, Zhuang Y, Lee P, Chi-Mei Li J, Wong PWC. Leisure and problem gaming behaviors among children and adolescents during school closures caused by covid-19 in hong kong: Quantitative cross-sectional survey study. *JMIR Serious Games.* 2021;9(2). doi: 10.2196/26808.
100. Donati MA, Guido CA, De Meo G, et al. Gaming among Children and Adolescents during the COVID-19 Lockdown: The Role of Parents in Time Spent on Video Games and Gaming Disorder Symptoms. *International Journal of Environmental Research and Public Health.* 2021;18(12).
101. Sheikh MK, Malavde R, Daigavane S. Yogic eye exercises followed by the ergonomic advice on eye fatigue in children attending online classes in COVID-19. *Intern J Cur Res Rev.* 2020;12(17):132-136. doi: 10.31782/IJCRR.2020.121720.
102. Coles-Brennan C, Sulley A, Young G. Management of digital eye strain. *Clin Exp Optom.* 2019;102(1):18-29. doi: 10.1111/cxo.12798.
103. Daum KM, Good G, Tijerina L. Symptoms in video display terminal operators and the presence of small refractive errors. *J Am Optom Assoc.* 1988;59(9):691-697.
104. Portello JK, Rosenfield M, Chu CA. Blink rate, incomplete blinks and computer vision syndrome. *Optom Vis Sci.* 2013;90(5):482-487. doi: 10.1097/OPX.0b013e31828f09a7.

105. Kawashima M, Sano K, Takechi S, Tsubota K. Impact of lifestyle intervention on dry eye disease in office workers: a randomized controlled trial. *Journal of Occupational Health*. 2018;60(4):281-288. doi: <https://doi.org/10.1539/joh.2017-0191-OA>.
106. Bhargava R, Kumar P, Phogat H, Kaur A, Kumar M. Oral omega-3 fatty acids treatment in computer vision syndrome related dry eye. *Contact Lens and Anterior Eye*. 2015;38(3):206-210. doi: 10.1016/j.clae.2015.01.007.
107. Deinema LA, Vingrys AJ, Wong CY, Jackson DC, Chinnery HR, Downie LE. A Randomized, Double-Masked, Placebo-Controlled Clinical Trial of Two Forms of Omega-3 Supplements for Treating Dry Eye Disease. *Ophthalmology*. 2017;124(1):43-52. doi: 10.1016/j.ophtha.2016.09.023.
108. Nosch DS, Foppa C, Tóth M, Joos RE. Blink Animation Software to Improve Blinking and Dry Eye Symptoms. *Optometry Vision Sci*. 2015;92(9).
109. World Health Organisation. *Guidelines on physical activity, sedentary behaviour and sleep for children under 5 years of age*. Geneva: ; 2019.
110. COUNCIL ON COMMUNICATIONS AND MEDIA. Media Use in School-Aged Children and Adolescents. *Pediatrics*. 2016;138(5):e20162592. doi: 10.1542/peds.2016-2592.
111. Simons M, Brug J, Chinapaw MJ, de Boer M, Seidell J, de Vet E. Replacing Non-Active Video Gaming by Active Video Gaming to Prevent Excessive Weight Gain in Adolescents. *PLoS One*. 2015;10(7):e0126023. doi: 10.1371/journal.pone.0126023 [doi].
112. Niermann CYN, Spengler S, Gubbels JS. Physical Activity, Screen Time, and Dietary Intake in Families: A Cluster-Analysis With Mother-Father-Child Triads. *Front Public Health*. 2018;6:276. doi: 10.3389/fpubh.2018.00276 [doi].
113. Faulkner G, Bassett-Gunter R, White L, Berry TR, Tremblay MS. Can The Moblees™ Move Canadian Children? Investigating the Impact of a Television Program on Children's Physical Activity. *Front Public Health*. 2018;6:206. doi: 10.3389/fpubh.2018.00206 [doi].

114. De Decker E, De Craemer M, De Bourdeaudhuij I, et al. Influencing factors of screen time in preschool children: An exploration of parents' perceptions through focus groups in six European countries. *Obes Rev.* 2012;13(SUPPL. 1):75-84. doi: 10.1111/j.1467-789X.2011.00961.x.
115. Xu H, Wen LM, Rissel C. Associations of parental influences with physical activity and screen time among young children: a systematic review. *J Obes.* 2015;2015:546925. doi: 10.1155/2015/546925.
116. Belmon LS, van Stralen MM, Busch V, Hamsen IA, Chinapaw MJM. What are the determinants of children's sleep behavior? A systematic review of longitudinal studies. *Sleep Med Rev.* 2019;43:60-70. doi: 10.1016/j.smr.2018.09.007.
117. Moon JH, Kim KW, Moon NJ. Smartphone use is a risk factor for pediatric dry eye disease according to region and age: a case control study. *BMC Ophthalmology.* 2016;16(1):188. doi: 10.1186/s12886-016-0364-4.