



**QUEEN'S
UNIVERSITY
BELFAST**

Diversity gradients of terrestrial vertebrates – substantial variations about a common theme

Raz, T., Allison, A., Avila, L. J., Bauer, A. M., Böhm, M., Caetano, G. H. D. O., Colli, G., Doan, T. M., Doughty, P., Grismer, L., Itescu, Y., Kraus, F., Martins, M., Morando, M., Murali, G., Nagy, Z. T., Nogueira, C. D. C., Novosolov, M., Oliver, P. M., ... Meiri, S. (2024). Diversity gradients of terrestrial vertebrates – substantial variations about a common theme. *Journal of Zoology*, 322(2), 126-140. <https://doi.org/10.1111/jzo.13130>

Published in:
Journal of Zoology

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2024 the authors.

This is an open access article published under a Creative Commons Attribution-NonCommercial-NoDerivs License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy























The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

ORIGINAL RESEARCH

Diversity gradients of terrestrial vertebrates – substantial variations about a common theme

T. Raz¹ , A. Allison² , L. J. Avila³ , A. M. Bauer⁴ , M. Böhm⁵ , G. H. de O. Caetano⁶, G. Colli⁷, T. M. Doan⁸ , P. Doughty⁹, L. Grismer¹⁰, Y. Itescu^{11,12,13} , F. Kraus¹⁴ , M. Martins¹⁵ , M. Morando^{3,16} , G. Murali⁶, Z. T. Nagy¹⁷ , C. de C. Nogueira¹⁸ , M. Novosolov¹⁹, P. M. Oliver^{20,21} , P. Passos²² , D. Pincheira-Donoso²³ , R. Sindaco²⁴ , A. Slavenko²⁵ , O. Torres-Carvajal²⁶ , P. Uetz²⁷ , P. Wagner^{4,28} , A. Zimin¹ , U. Roll^{6*} & S. Meiri^{1,29*} 

¹School of Zoology, Tel Aviv University, Tel Aviv, Israel

²Bishop Museum, Honolulu, HI, USA

³Grupo Herpetología Patagónica (GHP-LASIBIBE), Instituto Patagónico para el Estudio de los Ecosistemas Continentales (IPEEC), CONICET, Puerto Madryn, Argentina

⁴Department of Biology and Center for Biodiversity and Ecosystem Stewardship, Villanova University, Villanova, PA, USA

⁵Global Center for Species Survival, Indianapolis Zoo, Indianapolis, IN, USA

⁶Mitrani Department of Desert Ecology, The Jacob Blaustein Institutes of Desert Research, Ben-Gurion University of the Negev, Midreshet Ben-Gurion, Sede Boker, Israel

⁷Departamento de Zoologia, Instituto de Ciências Biológicas, Universidade de Brasília, Brasília, Brazil

⁸Division of Natural Sciences, New College of Florida, Sarasota, FL, USA

⁹Collections & Research, Western Australian Museum, Welshpool, WA, Australia

¹⁰Department of Biology, La Sierra University, Riverside, CA, USA

¹¹Leibniz Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

¹²Institute of Biology, Freie Universität Berlin, Berlin, Germany

¹³Department of Evolutionary and Environmental Biology, University of Haifa, Haifa, Israel

¹⁴Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA

¹⁵Departamento de Ecologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, Brazil

¹⁶Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB), Puerto Madryn, Argentina

¹⁷Independent Researcher, Berlin, Germany

¹⁸Departamento de Ecologia, Universidade de São Paulo, São Paulo, Brazil

¹⁹Lundbeck Foundation Genetics Center, GLOBE Institute, University of Copenhagen, Copenhagen, Denmark

²⁰Centre for Planetary Health and Food Security, Griffith University, Brisbane, QLD, Australia

²¹Biodiversity and Geosciences Program, Queensland Museum, South Brisbane, QLD, Australia

²²Departamento de Vertebrados, Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

²³School of Biological Sciences, Queen's University Belfast, Belfast, UK

²⁴Museo Civico di Storia Naturale, Carmagnola, Italy

²⁵Cesar Australia, Brunswick, VIC, Australia

²⁶Museo de Zoología, Escuela de Ciencias Biológicas, Pontificia Universidad Católica del Ecuador, Quito, Ecuador

²⁷Center for Biological Data Science, Virginia Commonwealth University, Richmond, VA, USA

²⁸Allwetterzoo, Münster, Germany

²⁹Steinhardt Museum of Natural History, Tel Aviv University, Tel Aviv, Israel

Keywords

biogeographical realms; environment–richness relationship; latitudinal diversity gradient; species richness patterns; terrestrial vertebrates; species richness; tetrapods; global-scaled research.

Correspondence

Tal Raz, School of Zoology, Tel Aviv University, Tel Aviv 6997801, Israel.
Email: raz@mail.tau.ac.il

Abstract

Environmental factors, such as temperature, precipitation, and elevation, explain most of the variation in species richness at the global scale. Nevertheless, richness patterns may have different drivers across taxa and regions. To date, a comprehensive global examination of how various factors such as climate or topography drive patterns of species richness across all terrestrial vertebrates, using the same methods and predictors, has been lacking. Recent advances in species-distribution data allowed us to model and examine the richness pattern of all terrestrial tetrapods comprehensively. We tested the relationship between environmental and biogeographical variables and richness of amphibians (5983 species), birds (9630), mammals (5004), reptiles (8939), and tetrapods as a whole, globally, and across biogeographical realms. We studied the effects of climatic, ecological, and

*Contributed equally to this work

Editor: Andrew Kitchener
Associate Editor: Kevin Arbuckle

Received 8 May 2023; revised 25 October 2023;
accepted 2 November 2023

doi:10.1111/jzo.13130

Introduction

The latitudinal diversity gradient, describing the increase in species richness from high to low latitudes, is one of the strongest and best-studied biogeographic patterns (Hillebrand, 2004; Lomolino *et al.*, 2017). It has been shown to prevail globally across taxa (e.g. plants – Kreft & Jetz, 2007; Scheiner & Rey-Benayas, 1994; invertebrates – Economo *et al.*, 2018, 2019; Pinkert *et al.*, 2022; vertebrates – Jetz & Fine, 2012; Wiens, 2007). Exceptions to the pattern are usually observed at lower spatial scales (i.e. locally, e.g. North America; Gaucherel *et al.*, 2018) and taxonomic scales (i.e. clades/taxa with relatively few species, e.g. pinnipeds; Cerezer *et al.*, 2022; Gaston, 1996).

Many hypotheses have been proposed to explain the latitudinal diversity gradient (Pontarp *et al.*, 2019). Hypotheses based on environmental drivers are usually strongly supported (Field *et al.*, 2009; Hawkins *et al.*, 2012; Hortal *et al.*, 2008). Climatic variables are thought to influence richness through various pathways (O'Brien, 1998, 2006; Storch *et al.*, 2012). According to the 'more-individuals hypothesis', resource availability, driven largely by temperature-water dynamics, limits the number of individuals and, consequently, the number of species (Srivastava & Lawton, 1998; Storch *et al.*, 2018; Wright, 1983). The 'evolutionary-rates hypothesis' proposes that diversification rates are faster in hot and humid places, resulting in greater species richness (Rohde, 1992). However, the relationship between diversification rates and current species richness is not always straightforward and, in some cases, faster diversification rates were found in temperate species-poor regions (e.g. Cerezer *et al.*, 2022; Rabosky, 2009; Rabosky *et al.*, 2018; Schluter, 2016). An additional hypothesis suggests that the long and relatively undisturbed evolutionary history of the tropics resulted in accumulation of more species than in temperate regions (Mittelbach *et al.*, 2007). Yet, past climates have been shown to explain less of the variation in current richness than current climates (Hawkins & Porter, 2003; Liang & Meiri, 2023). Environmental heterogeneity is also thought to substantially impact species richness (e.g. Hortal *et al.*, 2009; Stein *et al.*, 2014). A greater variety of habitats or topographic structures (usually represented by elevational range) is believed to result in numerous ecological niches which, in turn, fosters species specialization (Hortal *et al.*, 2013; MacArthur, 1964).

biogeographic drivers using generalized additive models. Richness patterns and their environmental associations varied among taxa and realms. Overall precipitation was the predominant richness predictor. However, temperature was more important in realms where both cold and warm conditions exist. In the Indomalayan realm, elevational range was very important. Richness patterns of mammals, birds, and amphibians were strongly related to precipitation whereas reptile richness was mostly associated with temperature. Our results support the universal importance of precipitation but also suggest that future global-scaled research should incorporate other relevant variables other than climate, such as elevational range, to gain a better understanding of the richness–environment relationship. By doing so, we can further advance our knowledge of the complex relationships between biodiversity and the environment.

Insularity is postulated to have an impact on species richness as islands typically harbor fewer species per unit area than the mainland (e.g. Ding *et al.*, 2006; Field *et al.*, 2009; Whittaker & Fernández-Palacios, 2007). Islands are associated with higher rates of extinction due to their small size and isolation, and lower rates of immigration (Foufopoulos & Ives, 1999; MacArthur & Wilson, 1967).

Richness interacts differently with environmental and ecological predictors across regions (e.g. Bohdalková *et al.*, 2021; Qian, 2009). Productivity-related factors explained most of the global variation in vertebrate richness (e.g. mammals – Barreto *et al.*, 2019; birds – Davies *et al.*, 2007; Hawkins *et al.*, 2007; amphibians – Gouveia *et al.*, 2013, mammals, amphibians, and birds – Bohdalková *et al.*, 2021; all classes – Barreto *et al.*, 2021). Productivity and water mostly affect richness in warm regions (e.g. Bohdalková *et al.*, 2021; Hawkins, Porter, *et al.*, 2003). Temperature is often weakly, and sometimes negatively, correlated with richness in warm regions, but is strongly and positively correlated with richness in cold (mainly northern) areas (e.g. Hawkins, Field, *et al.*, 2003; Kalmar & Currie, 2006). Elevational range is often strongly correlated with species richness in less seasonal areas, such as the tropics (e.g. Lewin *et al.*, 2016; Rahbek & Graves, 2001; Tallwin *et al.*, 2017), whereas the correlation is weaker in more seasonal regions (e.g. Qian *et al.*, 2007; Rodríguez *et al.*, 2005; Terribile & Diniz-Filho, 2009).

Due to this non-stationarity, some studies compare richness–environment relationships between geographical regions (e.g. Alves *et al.*, 2018; Barreto *et al.*, 2019, 2021), such as biogeographical realms. Realms are often used to geographically divide richness–environment models (Qian, 2009; Roll *et al.*, 2015; Voskamp *et al.*, 2017), or as an explanatory variable (Buckley & Jetz, 2007; Hawkins, Porter, *et al.*, 2003). At a coarse geographic scale, realms represent distinct species pools generated by mostly independent evolutionary histories (Falaschi *et al.*, 2023; Ricklefs, 2004; Smith *et al.*, 2005). Differences in diversification and dispersal between realms can be viewed as largely independent events of biodiversity accumulation, making realms obvious units for biogeographic analyses.

Global species-richness patterns of amphibians, birds, and mammals are largely congruent (Grenyer *et al.*, 2006). However, there are some distinctions among taxa, mostly at

regional scales—and when reptiles are compared to other tetrapod taxa (e.g. Currie, 1991; Hawkins *et al.*, 2012; Powney *et al.*, 2010; Roll *et al.*, 2017). Consequently, richness–environment relationships vary across taxa (Barreto *et al.*, 2021; Currie, 1991). Most notably, reptile richness pattern is commonly found to be closely associated with temperature, whereas bird, mammal, and amphibian richness are consistently correlated with water-related factors or with primary productivity (e.g. Barreto *et al.*, 2021; Hawkins, Field, *et al.*, 2003; Qian, 2010; Rodríguez *et al.*, 2005).

Most studies that examined richness–environment relationship of terrestrial vertebrates were confined to one or two tetrapod classes (Allen *et al.*, 2002; Araújo *et al.*, 2008; Barreto *et al.*, 2019; Costa *et al.*, 2007; Evans *et al.*, 2005; Foody, 2004; Fritz *et al.*, 2016; Hawkins & Porter, 2003; Kerr & Packer, 1997; Qian *et al.*, 2007; Rahbek & Graves, 2001; Rodríguez *et al.*, 2005), or involve birds, mammals, and amphibians (Belmaker & Jetz, 2011; Bohdalková *et al.*, 2021; Buckley & Jetz, 2007; Davies *et al.*, 2007; Gouveia *et al.*, 2013; Grenyer *et al.*, 2006; Gudex-Cross *et al.*, 2022; Hawkins *et al.*, 2007, 2012; Hortal *et al.*, 2008; O'Malley *et al.*, 2023; Wu & Liang, 2018). Studies that incorporate all tetrapods (including reptiles) have usually been confined to one region (Currie, 1991; Lewin *et al.*, 2016; Powney *et al.*, 2010; Tallowin *et al.*, 2017; Whittaker *et al.*, 2007).

Until recently, data on the distribution of tetrapods, particularly that of reptiles, were limited and lacked comprehensive coverage of species (Roll *et al.*, 2017). Earlier work, thus, relied on incomplete species datasets (Jetz & Fine, 2012; Qian, 2009, 2010; Qian & Ricklefs, 2008). Although some recent studies use the now available comprehensive datasets, they tend to focus on limited sets of environmental factors, primarily productivity, precipitation, and temperature (Marin *et al.*, 2018 at the bioregion level; Barreto *et al.*, 2021 at the grid-cell level). Other factors, which are known to be linked to species richness, such as environmental heterogeneity and insularity, have so far only been evaluated at local scales or using incomplete tetrapod-richness datasets (e.g. Hortal *et al.*, 2009; Howard *et al.*, 2020; Stein *et al.*, 2014; Tallowin *et al.*, 2017). Although these factors are generally considered less important than climate (e.g. Barreto *et al.*, 2019; Belmaker & Jetz, 2011), incorporating them into a broad spatial and taxonomic analysis could offer new and intriguing insights.

We suggest that the comprehensive species-distribution data that have recently become available for all tetrapod taxa—most recently for reptiles (Caetano *et al.*, 2022; Roll *et al.*, 2017)—allow to accurately test their global and regional richness patterns and their correlations to various environmental parameters. We examined the global richness patterns of all amphibians, reptiles, birds, and mammals together and tested their relationships with environmental, topographic, insularity, and biogeographic variables. We further modeled reptile, amphibian, bird, and mammal richness separately, with common scales, model features, and explanatory variables, to enable meaningful comparisons between them. We compared the results within biogeographical realms presenting a comprehensive global analysis of the effects of varied environmental attributes on gamma diversity.

Materials and methods

We obtained distributional data for amphibians and mammals from the IUCN (2021), for birds from the BirdLife International data zone (BirdLife International & NatureServe, 2019), and for reptiles from an updated version of Roll *et al.* (2017) (GARD 1.7, Caetano *et al.*, 2022). Data were filtered to include only native distributions of species and their breeding ranges (where designated as such). We created species-richness maps at a 96×96 km resolution (using a Behrmann equal-area projection, roughly 1×1 degree at the equator). We excluded cells with land comprising $<90\%$ of the entire area of the cell (e.g. coastal cells), including most small islands (Fig. 1). This procedure retains 5983 amphibian, 5004 mammal, 9630 bird, and 8939 reptile species (Table S1). Cells with no species were retained in the analyses, but Antarctica was excluded. While ensuring model integrity by considering cells with similar sizes is important, we acknowledge that including only cells with $<90\%$ land area may have downplayed the importance of insularity (see below). Therefore, we conducted an additional analysis incorporating all cells and including log-transformed area as an additional predictor. Preliminary analyses which incorporated cells with $>30\%$ and $>10\%$ land area yielded similar results to those of the original analyses, which included cells with $>90\%$ land area (results not shown).

To assess the similarity of global richness patterns among the four tetrapod classes, we computed correlation coefficients between the richness values of each pair of tetrapod classes. We used Pearson's correlation adjusted for spatial autocorrelation, using the 'Spatialpack' package (Osorio *et al.*, 2016).

We modeled richness using: (1) mean annual temperature (hereafter 'temperature', °C, 30-s resolution; from Karger *et al.*, 2017); (2) log of mean annual precipitation (hereafter 'precipitation', mm, 30-s resolution; Karger *et al.*, 2017). Because similar linear differences in precipitations are more meaningful in areas with relatively low precipitation (e.g. in the desert edge) than in high ones (e.g. in a rainforest; Egozcue *et al.*, 2006), we used a multiplicative (log) scale; (3) elevational range representing habitat heterogeneity. We downloaded mean elevation data (in m, 30-s resolution, from WorldClim 2.1 initiative, Fick & Hijmans, 2017) and calculated the elevational range in each 1×1 degree cell (see richness-map creation above) by subtracting the minimum elevation from the maximum elevation. We standardized these three factors to provide comparable regression coefficients; (4) the number of ecoregions in each grid cell (Dinerstein *et al.*, 2017) a coarse proxy of habitat diversity; and (5) insularity was treated as a categorical factor distinguishing between mainland and islands (Field *et al.*, 2009). We classified each cell according to whether it is located on the mainland or on an island (see Fig. S1 for the classification; the largest island is Greenland— ~ 2 mil km² and the smallest mainland is Australia— ~ 7.7 mil km²). Cells comprising both mainland and island regions were considered mainland. We deleted the following predictors that have variance-inflation factors >5 (VIF; Rogerson, 2006), and concurrencies higher than 0.8 (i.e. the presence of covariates that are themselves well modeled as




Figure 1 Species-richness patterns and latitudinal diversity gradient of terrestrial tetrapods. Richness of: (a) all tetrapods (amphibians, reptiles, birds, and mammals), (c) reptiles, (d) amphibians, (e) birds, (f) mammals. Light gray areas denote cells excluded from analyses because their land areas cover less than 90% of the cell's entire area. White land areas denote cells with richness values of zero. Dark blue colors denote regions with few species and red ones denote regions with many species (note that the scale differs between panels). Black lines denote borders between different realms. All maps are in an equal area Behrmann projection at a 96×96 km resolution. (b) Latitudinal diversity gradient depicted the change in global species richness along a latitudinal gradient. Regression lines of the relationship between latitude and richness were created using GAM with K parameter equals 10. For each group, the species-richness values were divided by their total number of species in order to depict relative species richness. Richness of all tetrapods together is depicted in black, reptiles in orange, amphibians in green, birds in blue, and mammals in pink.

smooth functions of other covariates in general additive models; Ito et al., 2005): (1) annual temperature range, (2) mean diurnal temperature range, (3) precipitation seasonality (all from Karger et al., 2017), (4) mean elevation, (5) habitat

homogeneity (based on NDVI; Tuanmu & Jetz, 2015), (6) net primary productivity (Imhoff & Bounoua, 2006), (7) interannual variation in mean temperature, (8) and total precipitation (both derived from Copernicus Climate Change Service

(C3S, 2017), (9) temperatures at the last glacial maximum (LGM), and (10) climate change velocity since the LGM (Karger et al., 2017). We also excluded (11) geographical location (longitude and latitude interaction), which we aimed to use to account for spatial autocorrelation, due to high concurrency with mean annual temperature.

We conducted the analyses globally and within biogeographical realms (of Olson et al., 2001), excluding Antarctica. In global models, we added biogeographical realms as a predictor. This measure of evolutionary/geological history is somewhat crude compared to other historical variables but represents distinctive lineage pools generated through historical interactions with biogeographical barriers (e.g. Ficetola et al., 2017; Smith et al., 2005).

To determine whether to use productivity or precipitation as predictors, we compared the AIC scores of models using either productivity or precipitation—and the other chosen predictors: temperature, elevational range, number of ecoregions, insularity, and realms for the global model. Models of precipitation had consistently lower AIC scores than models using productivity and we therefore used precipitation throughout.

We used generalized additive models (GAM) in order to incorporate nonlinear trends of the predictors and easily portray predictor–response relationships (Hastie & Tibshirani, 1990; Larsen, 2015). We analyzed GAM models of amphibians, birds, mammals, reptiles, and all tetrapods. We ran all analyses globally, and within each of the six major biogeographical realms (Neotropic, Nearctic, Palearctic, Afrotropic, Indomalaya, and Australasia), resulting in 35 GAM models. In all our models, the response variables were species richness per grid cell. We employed a log identity link function in all models, with a negative binomial error structure, which best fitted the data distribution and avoided overdispersion. For the continuous variables, precipitation, temperature, and elevational range, we examined the inclusion of interactions between predictors by trying all possible combinations. The K parameter value, which sets the upper limit on the degrees of freedom associated with the smooth function of the model, was also examined and incorporated with an automatic smoothness selection (Wood, 2017). We added insularity and the number of ecoregions as linear predictors. We evaluated model performance by R^2 and AIC values. We conducted all analyses using R 4.0 (R Core Team, 2021). We conducted GAM analyses, and concurrency testing, using the ‘mgcv’ package (Wood, 2017), and VIF analyses using the ‘car’ package (Fox & Weisberg, 2019).

To evaluate predictor importance, we checked the percentage decrease in R^2 between the model including all variables, and the model excluding each predictor. This decrease represents the additional variance in species richness explained by the predictor and its interactions that is not explained by the others. We produced partial-dependence plots that show the marginal effect each predictor has on model predictions (Hastie et al., 2001).

Results

Global species-richness patterns

All tetrapod taxa show very similar global richness patterns (Fig. 1, Table 1) but with some notable exceptions. Tetrapod richness is highest near the Andes, especially along the Amazon basin, in northern Amazonia (Fig. 1). It is also high along southeastern South America, along the African Great Lakes, and through much of Southeast Asia. Richness is low at high latitudes and desert regions worldwide. Birds and mammals show very similar patterns to tetrapods combined (Table 1) except that mammals are species poor in Australia (Fig. 1). Amphibians are conspicuously absent from most of the world’s deserts but are relatively rich in southeastern United States and eastern Madagascar. Reptile richness is the least well correlated with tetrapods and the other taxa (Table 1). Reptile richness is relatively high in deserts globally and in Australia.

Latitudinal richness gradients

All taxa display strong latitudinal richness gradients. Richness peaks just south of the equator and decreases toward higher latitudes (Fig. 1b). Amphibians show the most marked tropical peak and the steepest latitudinal decline. Reptiles decline most slowly away from the tropics at mid-latitudes, especially in the southern hemisphere where richness is relatively high even at latitudes higher than 30°S. Other taxa (especially amphibians) decline more steeply. In the Northern Hemisphere, all taxa steeply decline away from the equator and reach a local minimum around the 25th parallel (roughly the southern Sahara but south of the Tibetan Plateau). Reptile richness, however, remains stable around the Tropic of Cancer where it is relatively higher than that of other taxa. At higher latitudes, reptile, and amphibian richness decreases, while endotherm richness actually increases slowly until around 40°N (mammals) and even 50°N (birds), then decreases toward the North Pole. Birds and

Table 1 Correlations of species richness among tetrapods using Pearson’s correlation corrected for spatial autocorrelation

	d.f.					
	Taxon	Tetrapods	Reptiles	Amphibians	Birds	Mammals
R	Tetrapods		59.82	74.85	146.04	80.24
	Reptiles	0.86		59.74	68.22	66.31
	Amphibians	0.88	0.78		90.99	102.59
	Birds	0.98	0.76	0.82		90.99
	Mammals	0.96	0.78	0.83	0.93	

Correlation coefficients (R) in the bottom triangle. Degrees of freedom (d.f.) in the upper triangle. P -values are <0.0001 for all correlations.

mammals reach higher latitudes than reptiles and amphibians in the Northern, but not in the Southern Hemisphere (Antarctica excluded; Fig. 1).

Species-richness models

Chosen model parameters were consistent among all taxa and realms and set with temperature, precipitation, elevational range, and their interaction, with K parameter set at 150, and insularity and number of ecoregions as linear predictors. For global models, biogeographical realms were also set as a linear predictor. Environmental and biogeographical predictors explained 81% (amphibians), 82% (birds), 85% (mammals), 89% (all tetrapods), and 91% (reptiles) of the variation in global models (Table 2). For tetrapods and birds, all realm-specific models explained over 80% of the variation. For other taxa, models for the Afrotropical and Indomalayan realms were weaker ($R^2 = 58\text{--}74\%$). Models for Neotropical amphibians and Palearctic mammals had R^2 values of 73 and 76%, respectively. All other models had R^2 values exceeding 80%. Models of all taxa in the Nearctic were especially strong (R^2 values 94–96%), followed by models for the Australasian realm (88–92%, except for reptiles with 81%).

Predictor effects

Globally, precipitation was the most important predictor of richness patterns for all taxa except reptiles (Figs 2a and 4, Table S3), for which temperature was most important. The two predictors were positively associated with richness. Reptile richness peaked at maximum precipitation levels followed by amphibians that peaked at near maximum levels. However, the trend for reptile richness was weaker, with relatively high richness also seen at low precipitation levels. In contrast, the trend was very strong for amphibian richness, with extremely low richness at low precipitation levels. Endotherms showed an intermediate pattern (Fig. 2b). At low temperatures, ectotherms richness was lower compared to endotherms, while at the highest temperatures reptiles exhibited the highest richness among all taxa, resulting in the steepest trend. At the global scale, other predictors were less important. The richness of all taxa increased with the number of ecoregions and was lower on islands (Fig. 2c). Elevational range had a positive association with all groups at very low levels and a negative association at high levels (except for mammals). All groups' richness steeply increased up to ~750 m. Above that, mammal

richness slowly increased toward the highest elevational range levels. Reptile, tetrapod, and bird richness were fairly constant up to ~2500 m, above which reptile richness decreased toward the highest elevational ranges, while tetrapod and bird richness increased up to ~4500 m and then decreased. For amphibians, above ~750 m richness decreased up to ~2000 m then mildly increased toward ~5500 m, above which it declined.

Precipitation and temperature usually had the strongest (almost always positive) association with species richness across realms and taxa (Figs 3 and 4, Fig. S2, Table S3). Elevational range generally had a lower, more complex, association with richness, which varied across realms and taxa. However, its importance exceeded the one of precipitation or temperature in some cases (e.g. Indomalayan birds). Insularity (lower richness on islands than on continents in most realms and taxa) and the number of ecoregions (positively associated with richness) were usually the least important predictors (Fig. 3, Fig. S2).

Richness of different taxa was usually similarly related to the predictors within each realm, but differences exist (Figs 3 and 4, Fig. S2). Reptile richness was positively and strongly related to temperature, even within realms in which temperature had a weak or even a negative relationship with richness for the other taxa (most notably in the Australasian and Palearctic realms). Precipitation had a positive and strong association with amphibian richness in all realms, often stronger than for all other taxa (i.e. in the Nearctic, Afrotropic, and Palearctic realms; Figs 3 and 4, Fig. S2).

Predictors' association with richness varied between realms. Precipitation had an exceptionally strong positive association with Afrotropical taxa's richness whereas temperature had a weak association (Figs 3 and 4). In contrast, in the Nearctic, temperature had a very strong, generally positive association, whereas precipitation had a moderately negative relationship with tetrapod and endotherm richness, an extremely weak relationship with reptile richness, and a positive relationship with amphibian richness (Figs 3 and 4). In the Indomalayan realm, precipitation and elevational range were the most important predictors for the richness of all taxa (Fig. 4)—precipitation was positively associated with richness, whereas elevational range had a largely bimodal association. Richness peaked at ~2000 for all taxa, above which amphibian richness decreased while reptile and tetrapod richness peaked again and at ~5500 m and mammal and bird richness increased towards maximum elevational range levels (Fig. 3). In the Neotropics

Table 2 The adjusted R^2 of all the regression models explaining richness patterns across realms and for different taxa

Taxon/Realm	Neotropics	Afrotropics	Australasia	Indomalaya	Nearctic	Palearctic	Global
Amphibians	0.73	0.67	0.91	0.58	0.95	0.82	0.81
Birds	0.80	0.81	0.88	0.83	0.94	0.82	0.82
Mammals	0.87	0.74	0.92	0.66	0.95	0.76	0.85
Reptiles	0.91	0.67	0.81	0.71	0.96	0.85	0.91
Tetrapods	0.86	0.80	0.91	0.80	0.96	0.80	0.89

We built general additive models for each taxon, globally and within the different realms, resulting in 35 models. We modeled species richness as a function of environmental and biogeographical predictors.




Figure 2 Global associations of environmental and biogeographical predictors with tetrapod richness. Species richness within each group was rescaled to range between 0 (lowest richness) and 1 (highest richness) using the formula: $z_i = \frac{x_i - \min(x)}{\max(x) - \min(x)}$ | $i = 1, 2, \dots, n$. Where x_i represents the original data and z_i the rescaled data. (a) Predictors' importance in explaining richness variation; rows denote the different predictors used in the model. A predictor importance (ranges from 0 to 72) is the percentage decrease in R^2 between the model including all variables and the model excluding the predictor. (b, c) Partial dependence plots representing the change in rescaled predicted global species richness along variable gradients: (b) the two most important predictors of species richness of all classes, (c) predictors with lower importance. Global models were fitted for each taxon separately (reptiles depicted in orange, amphibians in green, birds in blue, mammals in pink, and all tetrapods together in black). Annual precipitation is on a logarithmic scale, but raw values are presented.

and Australasia, precipitation had the strongest association with richness of all taxa except reptiles which were most strongly associated with temperature (Fig. 4). In the Palearctic realm, temperature and precipitation dominated all models, with elevational range third in importance for most taxa. Additionally, for amphibian richness, precipitation was much more important than temperature, whereas the reverse was true for reptiles (for which elevational range importance exceeded the one of precipitation).

Models including all cells (as opposed to cells >90% land) had nearly consistently lower R^2 values (Table S2). However, insularity became an important negative predictor of richness in the Neotropics for tetrapods, birds, and mammals, and in Australasia for reptiles and amphibians (Table S4).

Discussion

Our study provides a comprehensive account of key environmental and biogeographical factors that underlie the richness patterns of all terrestrial vertebrates. We support previous findings (e.g. Barreto *et al.*, 2021; Qian, 2010) that precipitation is most strongly (and positively) related to tetrapod species richness (except reptiles), globally and largely in all realms except for the Nearctic (Figs 2–4). We show that associations between species richness and elevational range differ between taxa, globally and regionally, especially between endotherms and ectotherms (Figs 2 and 3). Moreover, we show that elevational range is a very strong predictor of species richness in Indomalaya while insularity has minor importance in explaining




Figure 3 The associations of environmental and biogeographical predictors with tetrapod richness within the different realms. Predictor (= column) relationship with rescaled richness, showing marginal effects on predicted species richness in each of the six biogeographical realms (rows). The marginal effect of each predictor is represented by a partial dependence plot. Models were fitted for each taxon separately (reptiles: orange, amphibians: green, birds: blue, mammals: pink, all tetrapods: black). Species richness of each group (y-axis) was scaled to range between 0 (lowest richness) and 1 (highest richness). Annual precipitation is on a logarithmic scale, but the raw values are presented.

variation in species richness (Fig. 4). Within realms, we find great variability in environment–richness relationships (consistent with Qian, 2010).

Taxonomic variation in richness–environment relationships

The overall tetrapod-richness pattern is very similar to that of birds and mammals (Table 1). This is probably because bird species have the largest ranges (Li et al., 2016) and amphibians and reptiles the smallest, thus, since large-ranged species influence more cells, endotherms more strongly influence tetrapod richness patterns (compare scales in Fig. 1; Jetz & Rahbek, 2002; Lennon et al., 2004; White et al., 2023). Consequently, the richness of birds, mammals, and tetrapods was similarly correlated with the environmental predictors, whereas amphibians and reptiles, often displayed different relationships (Figs 2–4; e.g. Barreto et al., 2021; Belmaker & Jetz, 2011; Buckley & Jetz, 2007; Jetz & Fine, 2012).

Endotherms are more tolerant of cold temperatures than ectotherms, and many birds avoid harsh seasonal environments through migration (Buckley et al., 2012; Hurlbert &

Haskell, 2003). However, endotherms require high and continuous food supply (Buckley et al., 2012; Pough, 1980; Shine, 2005). Thus, their richness is presumably less limited by solar energy than by resource availability. Accordingly, precipitation, which has considerable impacts on productivity (Liu et al., 2020), was overall the most important predictor of bird and mammal richness (Fig. 4). While consistent with most prior research (e.g. Hawkins, Field, et al., 2003; Rodríguez et al., 2005), Qian and Ricklefs (2008; at the country level, for all tetrapods) and Belmaker and Jetz (2011; for mammal assemblages) found that temperature indices were more important than water indices in predicting global species richness. It is worth noting that this could be due to the limited desert-region samples (refer to Fig. 1 in both papers), pivotal in global precipitation patterns. Moreover, data accessibility may have favored samples from Nearctic and Western European regions, known for their wide temperature ranges (Fig. S3; Bohdalková et al., 2021). Howard et al. (2020, Fig. S9) found that temperature seasonality was more important than precipitation for tetrapod, mammal, and bird species richness. However, it is worth considering that the inclusion of highly correlated factors, such as temperature seasonality and temperature, may




Figure 4 Spatial variance in the importance of environmental predictors on tetrapod species richness. The size of a circle is in accordance with its importance value calculated as the percentage decrease in R^2 between the model including all variables and the model excluding the predictor. The predictor importance is depicted for each realm, and the global model is presented at the bottom. The number of ecoregions is excluded due to its small impact on species richness in our models. Grouping of different taxa (e.g. mammals, birds, and tetrapods in the Afrotropics) occurs when the predictor importance is very similar between the different taxa. In such a case, the circle size corresponds to the mean of the importance values. The colored icons depict the environmental and biogeographical predictors used in the models (annual precipitation in blue, temperature in orange, elevation range in green, insularity in pink, and realms in gray).

have influenced their conclusions. Reptiles, which display a strong and positive relationship with temperature (Figs 2–4; in accordance with e.g. Barreto *et al.*, 2021; Qian & Ricklefs, 2008; Qian *et al.*, 2007), are highly dependent on ambient temperatures, but can cope with dry conditions (Buckley *et al.*, 2012; Schall & Pianka, 1978; e.g. see the high richness in the deserts of Australia in Fig. 1c). Amphibians, for which precipitation had the strongest positive correlation (Figs 3 and 4, Fig. S2; in accordance with, e.g. Buckley & Jetz, 2007), require readily available water or humid conditions for key physiological processes and activities (Feder & Burggren, 1992; Tracy, 1976).

Our research revealed that elevational range often exhibits distinct associations with the different taxa, often varying between endothermic and ectothermic species richness (Figs 2c and 3). Elevational range is thought to represent habitat heterogeneity and usually has a positive impact on species richness (e.g. Field *et al.*, 2009; Kerr & Packer, 1997; Stein *et al.*, 2014). For example, the peak in species richness around elevational range of ~4000 m in the Indomalayan and Palearctic realm (and to some extent in the global model; Figs 2c and 3) corresponds to the high habitat heterogeneity induced by the biogeographical barrier between the Indomalayan and the

Palearctic realm. However, elevational range is also correlated with mean elevation (global Pearson correlation coefficient 0.48). In very high elevations the assumption that greater elevational ranges include more available habitats is often not met due to harsher conditions (Hortal *et al.*, 2013). Thus, wide elevational ranges can negatively impact richness (e.g. Barreto *et al.*, 2019; Hortal *et al.*, 2013; Qian *et al.*, 2007). Accordingly, we found that for most regions and taxa, there is a decline in species richness at high values of elevational range (Figs 2c and 3). Moreover, ectotherms show in some regions (Global, Indomalaya, and Neotropical) a steeper or earlier (from lower elevational ranges) decline in species richness than endotherms (Fig. 3). It is possible that endotherms better cope with the cold conditions at high elevations (Buckley *et al.*, 2012).

Spatial variation in richness–environment relationships

Previous studies found that the relationship between environmental predictors and tetrapod richness differs across space (e.g. Alves *et al.*, 2018; Barreto *et al.*, 2021; Davies *et al.*, 2007). Bohdalková *et al.* (2021) suggested that the relationship between richness and temperature is only strong in

regions with a wide temperature range. In contrast, the relationship between richness and productivity appears to be more universal. Unlike temperature, our analysis suggests that high variation in elevational range does not necessarily lead to a strong correlation with richness (Fig. S3). Such a strong correlation is apparent in Indomalaya, but not in the Neotropics or Australasia, despite elevational range having a higher variation than precipitation and temperature in these regions. The large area of relatively homogeneous environment of the wet and warm Amazon, may harbor many large-ranged species and high range overlap, leading to higher richness (James & Shine, 2000), which is unrelated to the variation in elevational range (similarly, the hot and arid zone in Australia may induce a lower species richness for most taxa).

We found a negative relationship between temperature and richness in Australasia for amphibians, birds, and mammals and a humped-shape relationship in the Palearctic for these taxa (Fig. 3). Barreto *et al.* (2021) also found mixed trends in most realms, but for Nearctic birds, we found a richness peak at medium temperatures ($\sim 5^{\circ}\text{C}$) while Barreto *et al.* (2021) found a positive association. Our focus solely on breeding ranges, as opposed to their inclusion of non-breeding grounds in warmer regions, likely accounts for the temperature-dependent difference in species richness.

The latitudinal diversity gradient

The decrease in richness away from the equator is not symmetrical in any taxon (Fig. 1b). Species richness in the southern hemisphere is usually higher than in equivalent latitudes in the northern hemisphere in many taxa (e.g. Blackburn & Gaston, 1996; Burns, 2007; Dunn *et al.*, 2009). This might be explained by the higher temperatures in southern latitudes, which result from an extensive area of oceans in comparison with the more continental northern hemisphere (Chown *et al.*, 2004; Gaston & Chown, 1999).

Arid areas are much more extensive in the northern hemisphere than in the southern one. The Sahara Desert alone accounts for 45% of the world's hyperarid zones (FAO, 2019). This great desert (together with the Tibetan Plateau) potentially causes the steeper decline in richness from the equator toward high latitudes in the northern, versus the southern, hemisphere (Fig. 1b). North of these drylands, precipitation levels increase while temperatures decrease (Fig. S4), creating suitable conditions for many endotherm species, but less so for amphibians and reptiles (Fig. 1). The rise in tetrapod, bird, and mammal richness from 25° to 45° latitude contradicts the usual trend of diversity decreasing with latitude (Gaucherel *et al.*, 2018). Instead, it corresponds with higher levels of precipitation and productivity.

Caveats

Species-richness data are essential for understanding the history, ecology, and conservation priorities (Jenkins *et al.*, 2013). However, they suffer from data shortfalls. Most notably, there is a great spatial sampling bias, as diversity in deserts, high mountains, and geopolitically less stable, or poorer countries,

are less studied than, e.g. the USA, Canada, Australia, and Europe (e.g. Hickisch *et al.*, 2019; Marshall & Strine, 2019; Meyer *et al.*, 2015). Such biases reflect both the dearth of samples and often the lack of the necessary taxonomic expertise to properly identify the faunas of these regions. These Linnean and Wallacean shortfalls might cause inaccuracies in species-richness maps, which rely on global knowledge of species distributions (Hughes *et al.*, 2021; Yang *et al.*, 2013). However, as undersampling is probably strongest in the tropics, which have the highest richness values despite this bias, and our models had high explanatory power (Table 1), we think the main richness–environment patterns we found are generally robust. Nonetheless, spatial variation in biodiversity knowledge is significant even within the tropics (e.g. the Neotropics vs. Afrotropics and Indomalaya; Martin *et al.*, 2012; Wallace, 1859) which might more strongly impact our results. Additionally, many terrestrial vertebrates, particularly amphibians and reptiles, are still undescribed and are assumed to exist mainly in the tropics (e.g. Giam *et al.*, 2011; Melville *et al.*, 2021; Moura & Jetz, 2021). However, newly described species most likely have small range sizes (Giam *et al.*, 2011; Meiri, 2016), and thus their impact on large-scale species-richness patterns is probably small.

Conclusions

We have built upon recent advances in the availability of species-distribution data (Caetano *et al.*, 2022; IUCN, 2021; Roll *et al.*, 2017) and conducted thorough analyses of richness–environment relationships for terrestrial vertebrates, both collectively and separately for birds, mammals, amphibians, and reptiles at both global and biogeographical realms levels. While numerous studies have examined many of the patterns we present in this study, individually or partially, conducting them within a single study, while providing detailed richness–environment associations, allows us to make solid comparisons among variables, taxa, and regions. We support some earlier observations that precipitation is generally the most important factor to predict tetrapod species richness. Nevertheless, we also highlight two main exceptions: (1) reptile richness is strongly associated with temperature—as can be expected given reptiles' physiological and ecological needs, and (2) in the Nearctic, and to some extent the Palearctic, temperature is often the strongest predictor of richness—probably due to the high variance in temperatures across these realms. We further show that while elevational range is usually less important than climate, it is highly influential in the Indomalaya, and its relationship with species richness differs among taxa, especially between endotherms and ectotherms. Recent studies on species richness extensively investigated high-resolution richness–environment relationships and provided interesting insights (Barreto *et al.*, 2021; Bohdalková *et al.*, 2021). However, their focus has been limited to climatic variables. We suggest that future studies consider incorporating other relevant variables such as elevational range into their analyses. Such investigations may yield further novel insights and advance our understanding of the complex relationships between biodiversity and the environment.

Acknowledgments

This work was supported by the Israel Science Foundation (ISF; grant no. 406/19). We heartily thank Dr. Reut Vardi for valuable insights and discussions on the paper.

Author contributions

TR, SM, and UR conceived the ideas and designed the study; All authors collected the data; TR and UR analyzed the data; TR led the writing of the paper; All authors contributed critically to the drafts and gave final approval for publication.

Data availability statement

The distributional data used in this study to create the species richness maps can be obtained from the mentioned sources in the text.

References

- Allen, A. P., Brown, J. H., & Gillooly, J. F. (2002). Global biodiversity, biochemical kinetics, and the energetic-equivalence rule. *Science*, **297**, 1545–1548.
- Alves, D. M. C. C., Diniz-Filho, J. A. F., da Silva e Souza, K., Gouveia, S. F., & Villalobos, F. (2018). Geographic variation in the relationship between large-scale environmental determinants and bat species richness. *Basic and Applied Ecology*, **27**, 1–8.
- Araújo, M. B., Nogués-Bravo, D., Diniz-Filho, J. A. F., Haywood, A. M., Valdes, P. J., & Rahbek, C. (2008). Quaternary climate changes explain diversity among reptiles and amphibians. *Ecography*, **31**, 8–15.
- Barreto, E., Graham, C. H., & Rangel, T. F. (2019). Environmental factors explain the spatial mismatches between species richness and phylogenetic diversity of terrestrial mammals. *Global Ecology and Biogeography*, **28**, 1855–1865.
- Barreto, E., Rangel, T. F., Coelho, M. T. P., Cassemiro, F. A. S., Zimmermann, N. E., & Graham, C. H. (2021). Spatial variation in direct and indirect effects of climate and productivity on species richness of terrestrial tetrapods. *Global Ecology and Biogeography*, **30**, 1899–1908.
- Belmaker, J., & Jetz, W. (2011). Cross-scale variation in species richness–environment associations. *Global Ecology and Biogeography*, **20**, 464–474.
- BirdLife International, & NatureServe. (2019). *Bird species distribution maps of the world, version 5.0*. BirdLife International and NatureServe.
- Blackburn, T. M., & Gaston, K. J. (1996). Spatial patterns in the species richness of birds in the New World. *Ecography*, **19**, 369–376.
- Bohdalková, E., Toszogyova, A., Šímová, I., & Storch, D. (2021). Universality in biodiversity patterns: Variation in species–temperature and species–productivity relationships reveals a prominent role of productivity in diversity gradients. *Ecography*, **44**, 1366–1378.
- Buckley, L. B., Hurlbert, A. H., & Jetz, W. (2012). Broad-scale ecological implications of ectothermy and endothermy in changing environments. *Global Ecology and Biogeography*, **21**, 873–885.
- Buckley, L. B., & Jetz, W. (2007). Environmental and historical constraints on global patterns of amphibian richness. *Proceedings of the Royal Society B: Biological Sciences*, **274**, 1167–1173.
- Burns, K. C. (2007). Is tree diversity different in the Southern Hemisphere? *Journal of Vegetation Science*, **18**, 307–312.
- Caetano, G. H. O., Chapple, D. G., Grenyer, R., Raz, T., Rosenblatt, J., Tingley, R., Böhm, M., Meiri, S., & Roll, U. (2022). Automated assessment reveals that the extinction risk of reptiles is widely underestimated across space and phylogeny. *PLoS Biology*, **20**, e3001544.
- Cerezer, F. O., Machac, A., Rangel, T. F., & Dambros, C. S. (2022). Exceptions to the rule: Relative roles of time, diversification rates and regional energy in shaping the inverse latitudinal diversity gradient. *Global Ecology and Biogeography*, **31**, 1794–1809.
- Chown, S. L., Sinclair, B. J., Leinaas, H. P., & Gaston, K. J. (2004). Hemispheric asymmetries in biodiversity—A serious matter for ecology. *PLoS Biology*, **2**, e406.
- Copernicus Climate Change Service (C3S). (2017). *ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate*. Copernicus Climate Change Service Climate Data Store (CDS). <https://cds.climate.copernicus.eu/cdsapp#!/home>
- Costa, G. C., Nogueira, C., Machado, R. B., & Colli, G. R. (2007). Squamate richness in the Brazilian Cerrado and its environmental–climatic associations. *Diversity and Distributions*, **13**, 714–724.
- Currie, D. J. (1991). Energy and large-scale patterns of animal- and plant-species richness. *The American Naturalist*, **137**, 27–49.
- Davies, R. G., Orme, C. D. L., Storch, D., Olson, V. A., Thomas, G. H., Ross, S. G., Ding, T. S., Rasmussen, P. C., Bennett, P. M., Owens, I. P. F., Blackburn, T. M., & Gaston, K. J. (2007). Topography, energy and the global distribution of bird species richness. *Proceedings of the Royal Society B: Biological Sciences*, **274**, 1189–1197.
- Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E. C., Jones, B., Barber, C. V., Hayes, R., Kormos, C., Martin, V., Crist, E., ... Saleem, M. (2017). An ecoregion-based approach to protecting half the terrestrial realm. *BioScience*, **67**, 534–545.
- Ding, T.-S., Yuan, H.-W., Geng, S., Koh, C.-N., & Lee, P.-F. (2006). Macro-scale bird species richness patterns of the East Asian mainland and islands: Energy, area and isolation. *Journal of Biogeography*, **33**, 683–693.
- Dunn, R. R., Agosti, D., Andersen, A. N., Arnan, X., Bruhl, C. A., Cerdá, X., Ellison, A. M., Fisher, B. L., Fitzpatrick, M. C., Gibb, H., Gotelli, N. J., Gove, A. D., Guenard, B., Janda, M., Kaspari, M., Laurent, E. J., Lessard, J. P., Longino, J. T., Majer, J. D., ... Sanders, N. J. (2009). Climatic drivers of hemispheric asymmetry in global patterns of ant species richness. *Ecology Letters*, **12**, 324–333.

- Economio, E. P., Huang, J.-P., Fischer, G., Sarnat, E. M., Narula, N., Janda, M., Guénard, B., Longino, J. T., & Knowles, L. L. (2019). Evolution of the latitudinal diversity gradient in the hyperdiverse ant genus *Pheidole*. *Global Ecology and Biogeography*, **28**, 456–470.
- Economio, E. P., Narula, N., Friedman, N. R., Weiser, M. D., & Guénard, B. (2018). Macroecology and macroevolution of the latitudinal diversity gradient in ants. *Nature Communications*, **9**, 1–8.
- Egozcue, J. J., Pawlowsky-Glahn, V., Ortego, M. I., & Tolosana-Delgado, R. (2006). The effect of scale in daily precipitation hazard assessment. *Natural Hazards and Earth System Sciences*, **6**, 459–470.
- Evans, K. L., Greenwood, J. J. D., & Gaston, K. J. (2005). Dissecting the species–energy relationship. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 2155–2163.
- Falaschi, M., Marta, S., Lo Parrino, E., Roll, U., Meiri, S., & Ficetola, G. F. (2023). Global bioregions of reptiles confirm the consistency of bioregionalization processes across vertebrate clades. *Global Ecology and Biogeography*, **32**, 1272–1284.
- FAO. (2019). *Trees, forests and land use in drylands: The first global assessment: Full report*. FAO.
- Feder, M. E., & Burggren, W. W. (Eds.). (1992). *Environmental physiology of the amphibians*. University of Chicago Press.
- Ficetola, G. F., Mazel, F., & Thuiller, W. (2017). Global determinants of zoogeographical boundaries. *Nature Ecology & Evolution*, **1**, 1–7.
- Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology*, **37**, 4302–4315.
- Field, R., Hawkins, B. A., Cornell, H. V., Currie, D. J., Diniz-Filho, J. A. F., Guégan, J. F., Kaufman, D. M., Kerr, J. T., Mittelbach, G. G., Oberdorff, T., O'Brien, E. M., & Turner, J. R. G. (2009). Spatial species–richness gradients across scales: A meta-analysis. *Journal of Biogeography*, **36**, 132–147.
- Foody, G. M. (2004). Spatial nonstationarity and scale-dependency in the relationship between species richness and environmental determinants for the sub-Saharan endemic avifauna. *Global Ecology and Biogeography*, **13**, 315–320.
- Foufopoulos, J., & Ives, A. R. (1999). Reptile extinctions on land-bridge islands: Life-history attributes and vulnerability to extinction. *The American Naturalist*, **153**, 1–25.
- Fox, J., & Weisberg, S. (2019). *An R companion to applied regression* (3rd ed.). Sage.
- Fritz, S. A., Eronen, J. T., Schnitzler, J., Hof, C., Janis, C. M., Mulch, A., Böhning-Gaese, K., & Graham, C. H. (2016). Twenty-million-year relationship between mammalian diversity and primary productivity. *Proceedings of the National Academy of Sciences of the United States of America*, **113**, 10908–10913.
- Gaston, K. J. (1996). Biodiversity - latitudinal gradients. *Progress in Physical Geography: Earth and Environment*, **20**, 466–476.
- Gaston, K. J., & Chown, S. L. (1999). Why Rapoport's rule does not generalise. *Oikos*, **84**, 309–312.
- Gauchere, C., Tramier, C., Devictor, V., Svenning, J.-C., & Hély, C. (2018). Where and at which scales does the latitudinal diversity gradient fail? *Journal of Biogeography*, **45**, 1905–1916.
- Giam, X., Scheffers, B. R., Sodhi, N. S., Wilcove, D. S., Ceballos, G., & Ehrlich, P. R. (2011). Reservoirs of richness: Least disturbed tropical forests are centres of undescribed species diversity. *Proceedings of the Royal Society B: Biological Sciences*, **279**, 67–76.
- Gouveia, S. F., Hortal, J., Cassemiro, F. A. S., Rangel, T. F., & Diniz-Filho, J. A. F. (2013). Nonstationary effects of productivity, seasonality, and historical climate changes on global amphibian diversity. *Ecography*, **36**, 104–113.
- Grenyer, R., Orme, C. D. L., Jackson, S. F., Thomas, G. H., Davies, R. G., Davies, T. J., Jones, K. E., Olson, V. A., Ridgely, R. S., Rasmussen, P. C., Ding, T. S., Bennett, P. M., Blackburn, T. M., Gaston, K. J., Gittleman, J. L., & Owens, I. P. F. (2006). Global distribution and conservation of rare and threatened vertebrates. *Nature*, **444**, 93–96.
- Gudex-Cross, D., Zhu, L., Keyser, S. R., Zuckerberg, B., Pauli, J. N., Radeloff, V. C., & Fleishman, E. (2022). Winter conditions structure extratropical patterns of species richness of amphibians, birds and mammals globally. *Global Ecology and Biogeography*, **00**, 1–15.
- Hastie, T., Friedman, J., & Tibshirani, R. (2001). Model inference and averaging. In *The elements of statistical learning: Data mining, inference, and prediction* (pp. 225–256). Springer.
- Hastie, T. J., & Tibshirani, R. J. (1990). *Generalized additive models*. Routledge.
- Hawkins, B. A., Alexandre, J., Diniz-Filho, F., Jaramillo, C. A., & Soeller, S. A. (2007). Climate, niche conservatism, and the global bird diversity gradient. *The American Naturalist*, **170**, 16–27.
- Hawkins, B. A., Field, R., Cornell, H. V., Currie, D. J., Guégan, J. F., Kaufman, D. M., Kerr, J. T., Mittelbach, G. G., Oberdorff, T., O'Brien, E. M., Porter, E. E., & Turner, J. R. G. (2003). Energy, water, and broad-scale geographic patterns of species richness. *Ecology*, **84**, 3105–3117.
- Hawkins, B. A., McCain, C. M., Davies, T. J., Buckley, L. B., Anacker, B. L., Cornell, H. V., Damschen, E. I., Grytnes, J. A., Harrison, S., Holt, R. D., Kraft, N. J. B., & Stephens, P. R. (2012). Different evolutionary histories underlie congruent species richness gradients of birds and mammals. *Journal of Biogeography*, **39**, 825–841.
- Hawkins, B. A., & Porter, E. E. (2003). Relative influences of current and historical factors on mammal and bird diversity patterns in deglaciated North America. *Global Ecology and Biogeography*, **12**, 475–481.
- Hawkins, B. A., Porter, E. E., & Felizola Diniz-Filho, J. A. (2003). Productivity and history as predictors of the latitudinal diversity gradient of terrestrial birds. *Ecology*, **84**, 1608–1623.

- Hickisch, R., Hodgetts, T., Johnson, P. J., Sillero-Zubiri, C., Tockner, K., & Macdonald, D. W. (2019). Effects of publication bias on conservation planning. *Conservation Biology*, **33**, 1151–1163.
- Hillebrand, H. (2004). On the generality of the latitudinal diversity gradient. *The American Naturalist*, **163**, 192–211.
- Hortal, J., Carrascal, L. M., Triantis, K. A., Thebault, E., Meiri, S., & Sfenthourakis, S. (2013). Species richness can decrease with altitude but not with habitat diversity. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E2149–E2150.
- Hortal, J., Rodríguez, J., Nieto-Díaz, M., & Lobo, J. M. (2008). Regional and environmental effects on the species richness of mammal assemblages. *Journal of Biogeography*, **35**, 1202–1214.
- Hortal, J., Triantis, K. A., Meiri, S., Thébault, E., & Sfenthourakis, S. (2009). Island species richness increases with habitat diversity. *The American Naturalist*, **174**, E205–E217.
- Howard, C., Flather, C. H., & Stephens, P. A. (2020). A global assessment of the drivers of threatened terrestrial species richness. *Nature Communications*, **11**, 993.
- Hughes, A. C., Orr, M. C., Yang, Q., & Qiao, H. (2021). Effectively and accurately mapping global biodiversity patterns for different regions and taxa. *Global Ecology and Biogeography*, **30**, 1–14.
- Hurlbert, A. H., & Haskell, J. P. (2003). The effect of energy and seasonality on avian species richness and community composition. *The American Naturalist*, **161**, 83–97.
- Imhoff, M. L., & Bounoua, L. (2006). Exploring global patterns of net primary production carbon supply and demand using satellite observations and statistical data. *Journal of Geophysical Research-Atmospheres*, **111**, D22S12.
- Ito, K., De Leon, S. F., & Lippmann, M. (2005). Associations between ozone and daily mortality: Analysis and meta-analysis. *Epidemiology*, **16**, 446–457.
- IUCN. (2021). *International Union for Conservation of Nature*. IUCN. <https://www.iucn.org>
- James, C. D., & Shine, R. (2000). Why are there so many coexisting species of lizards in Australian deserts? *Oecologia*, **125**, 127–141.
- Jenkins, C. N., Pimm, S. L., & Joppa, L. N. (2013). Global patterns of terrestrial vertebrate diversity and conservation. *Proceedings of the National Academy of Sciences of the United States of America*, **110**, E2602–E2610.
- Jetz, W., & Fine, P. V. A. (2012). Global gradients in vertebrate diversity predicted by historical area-productivity dynamics and contemporary environment. *PLoS Biology*, **10**, e1001292.
- Jetz, W., & Rahbek, C. (2002). Geographic range size and determinants of avian species richness. *Science*, **297**, 1548–1551.
- Kalmar, A., & Currie, D. J. (2006). A global model of island biogeography. *Global Ecology and Biogeography*, **15**, 72–81.
- Karger, D. N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R. W., Zimmermann, N. E., Linder, H. P., & Kessler, M. (2017). Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, **4**, 170122.
- Kerr, J. T., & Packer, L. (1997). Habitat heterogeneity as a determinant of mammal species richness in high-energy regions. *Nature*, **385**, 252–254.
- Kreft, H., & Jetz, W. (2007). Global patterns and determinants of vascular plant diversity. *Proceedings of the National Academy of Sciences of the United States of America*, **104**, 5925–5930.
- Larsen, K. (2015). GAM: The predictive modeling silver bullet. *Multithreaded Stitch Fix*, **30**, 1–27.
- Lennon, J. J., Koleff, P., Greenwood, J. J. D., & Gaston, K. J. (2004). Contribution of rarity and commonness to patterns of species richness. *Ecology Letters*, **7**, 81–87.
- Lewin, A., Feldman, A., Bauer, A. M., Belmaker, J., Broadley, D. G., Chirio, L., Itescu, Y., LeBreton, M., Maza, E., Meirte, D., Nagy, Z. T., Novosolov, M., Roll, U., Tallowin, O., Trape, J. F., Vidan, E., & Meiri, S. (2016). Patterns of species richness, endemism and environmental gradients of African reptiles. *Journal of Biogeography*, **43**, 2380–2390.
- Li, Y., Li, X., Sandel, B., Blank, D., Liu, Z., Liu, X., & Yan, S. (2016). Climate and topography explain range sizes of terrestrial vertebrates. *Nature Climate Change*, **6**, 498–502.
- Liang, T., & Meiri, S. (2023). Lizard richness in mainland China is more strongly correlated with energy and climatic stability than with diversification rates. *Journal of Biogeography*. <https://doi.org/10.1111/jbi.14736>
- Liu, J., Ma, X., Duan, Z., Jiang, J., Reichstein, M., & Jung, M. (2020). Impact of temporal precipitation variability on ecosystem productivity. *Wiley Interdisciplinary Reviews Water*, **7**, e1481.
- Lomolino, M. V., Riddle, B. R., & Whittaker, R. J. (2017). *Biogeography*. Oxford University Press.
- MacArthur, R. H. (1964). Environmental factors affecting bird species diversity. *The American Naturalist*, **98**, 387–397.
- MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography*. Princeton University Press.
- Marin, J., Rapacciuolo, G., Costa, G. C., Graham, C. H., Brooks, T. M., Young, B. E., Radeloff, V. C., Behm, J. E., Helmus, M. R., & Hedges, S. B. (2018). Evolutionary time drives global tetrapod diversity. *Proceedings of the Royal Society B: Biological Sciences*, **285**, 20172378.
- Marshall, B. M., & Strine, C. T. (2019). Exploring snake occurrence records: Spatial biases and marginal gains from accessible social media. *PeerJ*, **2019**, e8059.
- Martin, L. J., Blossey, B., & Ellis, E. (2012). Mapping where ecologists work: Biases in the global distribution of terrestrial ecological observations. *Frontiers in Ecology and the Environment*, **10**, 195–201.
- Meiri, S. (2016). Small, rare and trendy: Traits and biogeography of lizards described in the 21st century. *Journal of Zoology*, **299**, 251–261.
- Melville, J., Chapple, D. G., Keogh, J. S., Sumner, J., Amey, A., Bowles, P., Brennan, I. G., Couper, P., Donnellan, S. C., Doughty, P., Edwards, D. L., Ellis, R. J., Esquerré, D., Fenker,

- J., Gardner, M. G., Georges, A., Haines, M. L., Hoskin, C. J., Hutchinson, M., ... Tingley, R. (2021). A return-on-investment approach for prioritization of rigorous taxonomic research needed to inform responses to the biodiversity crisis. *PLoS Biology*, **19**, e3001210.
- Meyer, C., Krefl, H., Guralnick, R., & Jetz, W. (2015). Global priorities for an effective information basis of biodiversity distributions. *Nature Communications*, **6**, 1–8.
- Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M. B., Harrison, S. P., Hurlbert, A. H., Knowlton, N., Lessios, H. A., McCain, C. M., McCune, A. R., McDade, L. A., McPeck, M. A., Near, T. J., Price, T. D., Ricklefs, R. E., Roy, K., Sax, D. F., ... Turelli, M. (2007). Evolution and the latitudinal diversity gradient: Speciation, extinction and biogeography. *Ecology Letters*, **10**, 315–331.
- Moura, M. R., & Jetz, W. (2021). Shortfalls and opportunities in terrestrial vertebrate species discovery. *Nature Ecology & Evolution*, **5**, 631–639.
- O'Brien, E. (1998). Water-energy dynamics, climate, and prediction of woody plant species richness: An interim general model. *Journal of Biogeography*, **25**, 379–398.
- O'Brien, E. M. (2006). Biological relativity to water-energy dynamics. *Journal of Biogeography*, **33**, 1868–1888.
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., & Kassem, K. R. (2001). Terrestrial ecoregions of the world: A new map of life on earth. *BioScience*, **51**, 933.
- O'Malley, C. P. B., Roberts, G. G., Mannion, P. D., Hackel, J., & Wang, Y. (2023). Coherence of terrestrial vertebrate species richness with external drivers across scales and taxonomic groups. *Global Ecology and Biogeography*, **32**, 1285–1301.
- Osorio, F., Vallejos, R., & Cuevas, F. (2016). SpatialPack: Computing the association between two spatial processes. *arXiv*. <https://doi.org/10.48550/arXiv.1611.05289>
- Pinkert, S., Barve, V., Guralnick, R., & Jetz, W. (2022). Global geographical and latitudinal variation in butterfly species richness captured through a comprehensive country-level occurrence database. *Global Ecology and Biogeography*, **31**, 830–839.
- Pontarp, M., Bunnefeld, L., Cabral, J. S., Etienne, R. S., Fritz, S. A., Gillespie, R., Graham, C. H., Hagen, O., Hartig, F., Huang, S., Jansson, R., Maliet, O., Münkemüller, T., Pellissier, L., Rangel, T. F., Storch, D., Wiegand, T., & Hurlbert, A. H. (2019). The latitudinal diversity gradient: Novel understanding through mechanistic eco-evolutionary models. *Trends in Ecology & Evolution*, **34**, 211–223.
- Pough, F. H. (1980). The advantages of ectothermy for tetrapods. *The American Naturalist*, **115**, 92–112.
- Powney, G. D., Grenyer, R., Orme, C. D. L., Owens, I. P. F., & Meiri, S. (2010). Hot, dry and different: Australian lizard richness is unlike that of mammals, amphibians and birds. *Global Ecology and Biogeography*, **19**, 386–396.
- Qian, H. (2009). Global tests of regional effect on species richness of vascular plants and terrestrial vertebrates. *Ecography*, **32**, 553–560.
- Qian, H. (2010). Environment-richness relationships for mammals, birds, reptiles, and amphibians at global and regional scales. *Ecological Research*, **25**, 629–637.
- Qian, H., & Ricklefs, R. E. (2008). Global concordance in diversity patterns of vascular plants and terrestrial vertebrates. *Ecology Letters*, **11**, 547–553.
- Qian, H., Wang, X., Wang, S., & Li, Y. (2007). Environmental determinants of amphibian and reptile species richness in China. *Ecography*, **30**, 471–482.
- R Core Team. (2021). *R: A language and environment for statistical computing*. Austria.
- Rabosky, D. L. (2009). Ecological limits and diversification rate: Alternative paradigms to explain the variation in species richness among clades and regions. *Ecology Letters*, **12**, 735–743.
- Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., Kaschner, K., Garilao, C., Near, T. J., Coll, M., & Alfaro, M. E. (2018). An inverse latitudinal gradient in speciation rate for marine fishes. *Nature*, **559**, 392–395.
- Rahbek, C., & Graves, G. R. (2001). Multiscale assessment of patterns of avian species richness. *Proceedings of the National Academy of Sciences of the United States of America*, **98**, 4534–4539.
- Ricklefs, R. E. (2004). A comprehensive framework for global patterns in biodiversity. *Ecology Letters*, **7**, 1–15.
- Rodríguez, M. Á., Belmontes, J. A., & Hawkins, B. A. (2005). Energy, water and large-scale patterns of reptile and amphibian species richness in Europe. *Acta Oecologica*, **28**, 65–70.
- Rogerson, P. A. (2006). *Statistical methods for geography: A student's guide* (2nd ed.). SAGE Publications Ltd.
- Rohde, K. (1992). Latitudinal gradients in species diversity: The search for the primary cause. *Oikos*, **65**, 514.
- Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A. M., Bernard, R., Böhm, M., Castro-Herrera, F., Chirio, L., Collen, B., Colli, G. R., Dabool, L., Das, I., Doan, T. M., Grismer, L. L., Hoogmoed, M., Itescu, Y., Kraus, F., Lebreton, M., ... Meiri, S. (2017). The global distribution of tetrapods reveals a need for targeted reptile conservation. *Nature Ecology & Evolution*, **1**, 1677–1682.
- Roll, U., Geffen, E., & Yom-Tov, Y. (2015). Linking vertebrate species richness to tree canopy height on a global scale. *Global Ecology and Biogeography*, **24**, 814–825.
- Schall, J. J., & Pianka, E. R. (1978). Geographical trends in numbers of species. *Science*, **201**, 679–686.
- Scheiner, S. M., & Rey-Benayas, J. M. (1994). Global patterns of plant diversity. *Evolutionary Ecology*, **8**, 331–347.
- Schluter, D. (2016). Speciation, ecological opportunity, and latitude (American Society of Naturalists Address). *The American Naturalist*, **187**, 1–18.
- Shine, R. (2005). Life-history evolution in reptiles. *Annual Review of Ecology, Evolution, and Systematics*, **36**, 23–46.

- Smith, S. A., Stephens, P. R., & Wiens, J. J. (2005). Replicate patterns of species richness, historical biogeography, and phylogeny in Holarctic treefrogs. *Evolution*, **59**, 2433–2450.
- Srivastava, D. S., & Lawton, J. H. (1998). Why more productive sites have more species: An experimental test of theory using tree-hole communities. *The American Naturalist*, **152**, 510–529.
- Stein, A., Gerstner, K., & Kreft, H. (2014). Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecology Letters*, **17**, 866–880.
- Storch, D., Bohdalková, E., & Okie, J. (2018). The more-individuals hypothesis revisited: The role of community abundance in species richness regulation and the productivity–diversity relationship. *Ecology Letters*, **21**, 920–937.
- Storch, D., Keil, P., & Jetz, W. (2012). Universal species–area and endemics–area relationships at continental scales. *Nature*, **488**, 78–81.
- Tallowin, O., Allison, A., Algar, A. C., Kraus, F., & Meiri, S. (2017). Papua New Guinea terrestrial-vertebrate richness: Elevation matters most for all except reptiles. *Journal of Biogeography*, **44**, 1734–1744.
- Terribile, L. C., & Diniz-Filho, J. A. F. (2009). Spatial patterns of species richness in New World coral snakes and the metabolic theory of ecology. *Acta Oecologica*, **35**, 163–173.
- Tracy, C. R. (1976). A model of the dynamic exchanges of water and energy between a terrestrial amphibian and its environment. *Ecological Monographs*, **46**, 293–326.
- Tuanmu, M. N., & Jetz, W. (2015). A global, remote sensing-based characterization of terrestrial habitat heterogeneity for biodiversity and ecosystem modelling. *Global Ecology and Biogeography*, **24**, 1329–1339.
- Voskamp, A., Baker, D. J., Stephens, P. A., Valdes, P. J., & Willis, S. G. (2017). Global patterns in the divergence between phylogenetic diversity and species richness in terrestrial birds. *Journal of Biogeography*, **44**, 709–721.
- Wallace, A. R. (1859). The geographical distribution of birds. *Ibis*, **1**, 449–454.
- White, H. J., McKeon, C. M., Pakeman, R. J., & Buckley, Y. M. (2023). The contribution of geographically common and rare species to the spatial distribution of biodiversity. *Global Ecology and Biogeography*, **32**, 1730–1747.
- Whittaker, R. J., & Fernández-Palacios, J. M. (2007). *Island biogeography: Ecology, evolution, and conservation*. Oxford University Press.
- Whittaker, R. J., Nogués-Bravo, D., & Araújo, M. B. (2007). Geographical gradients of species richness: a test of the water-energy conjecture of Hawkins *et al.* (2003) using European data for five taxa. *Global Ecology and Biogeography*, **16**, 76–89.
- Wiens, J. J. (2007). Global patterns of diversification and species richness in amphibians. *The American Naturalist*, **170**, S86–S106.
- Wood, S. (2017). *Generalized additive models an introduction with R* (2nd ed.). Chapman and Hall/CRC.
- Wright, D. (1983). Species-energy theory: An extension of species-area theory. *Oikos*, **41**, 496–506.
- Wu, J., & Liang, S. (2018). Developing an integrated remote sensing based biodiversity index for predicting animal species richness. *Remote Sensing*, **10**, 739.
- Yang, W., Ma, K., & Kreft, H. (2013). Geographical sampling bias in a large distributional database and its effects on species richness-environment models. *Journal of Biogeography*, **40**, 1415–1426.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. The insularity predictor used in the models.

Figure S2. Predictors' importance for all regional models.

Figure S3. Spatial variation in the variance of environmental predictors and their importance on tetrapod species richness.

Figure S4. The continuous predictors used in the models.

Table S1. Species included in the creation of the species richness maps.

Table S2. The adjusted R^2 of all the regression models.

Table S3. Predictors' importance for all the models.

Table S4. Predictors' importance for all the models including all cells.