



**QUEEN'S  
UNIVERSITY  
BELFAST**

## Degradation of carbon disulphide (CS<sub>2</sub>) in soils and groundwater from a CS<sub>2</sub> contaminated site

Cox, S. F., McKinley, J. D., Ferguson, A. S., O'Sullivan, G., & Kalin, R. M. (2013). Degradation of carbon disulphide (CS<sub>2</sub>) in soils and groundwater from a CS<sub>2</sub> contaminated site. *Environmental Earth Sciences*, 68(7), 1935-1944. <https://doi.org/10.1007/s12665-012-1887-y>

**Published in:**  
Environmental Earth Sciences

**Document Version:**  
Peer reviewed version

**Queen's University Belfast - Research Portal:**  
[Link to publication record in Queen's University Belfast Research Portal](#)

### General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

### Open Access

This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

1  
2  
3  
4  
5  
6

The final publication is available at <http://link.springer.com/article/10.1007%2Fs12665-012-1881-y#page-1>

Cox, S. F., McKinley, J. D., Ferguson A. S., O'Sullivan, G. and Kalin, R. (2013) Degradation of carbon disulphide (CS<sub>2</sub>) in soils and groundwater from a CS<sub>2</sub> contaminated site, *Environ Earth Sci.* Volume 68, Issue 7, pp 1935-1944

7

## Degradation of carbon disulphide (CS<sub>2</sub>) in soils and

8

## groundwater from a CS<sub>2</sub> contaminated site

9

Siobhan F. Cox<sup>\*a</sup>, John D. McKinley<sup>a</sup>, Andrew S. Ferguson<sup>a1</sup>, Gwen O'Sullivan<sup>a</sup> and

10

Robert M. Kalin<sup>b</sup>

11

### *Affiliations*

12

<sup>a</sup> *Environmental Engineering Research Centre, School of Planning, Architecture and Civil Engineering, Queen's University Belfast, Belfast, BT9 5AG, United Kingdom*

13

14

Tel: +44 28 9097 4152

15

Fax: +44 28 9097 4278

16

Email: [s.cox@qub.ac.uk](mailto:s.cox@qub.ac.uk)

17

<sup>b</sup> *Department of Civil Engineering, University of Strathclyde, Glasgow, G1 1XN, Scotland, United Kingdom*

18

19

20

### *Current Addresses*

21

<sup>1</sup> *Civil Engineering and Engineering Mechanics, Columbia University, 605 Seeley W. Mudd, 500 West 120th Street, New York, NY 10027-6902, USA*

22

23

24

## **ABSTRACT**

25

This study is the first investigation of biodegradation of carbon disulphide (CS<sub>2</sub>) in soil that provides estimates

26

of degradation rates and identifies intermediate degradation products and carbon isotope signatures of

27

degradation. Microcosm studies were undertaken under anaerobic conditions using soil and groundwater

28

recovered from CS<sub>2</sub> contaminated sites. Proposed degradation mechanisms were validated using equilibrium

29

speciation modelling of concentrations and carbon isotope ratios. A first order degradation rate constant of

30

$1.25 \times 10^{-2} \text{ h}^{-1}$  was obtained for biological degradation with soil. Carbonyl sulphide (COS) and hydrogen

31

sulphide (H<sub>2</sub>S) were found to be intermediates of degradation, but did not accumulate in vials. A <sup>13</sup>C/<sup>12</sup>C

32

enrichment factor of  $-7.5 \pm 0.8\%$  was obtained for degradation within microcosms with both soil and

33

groundwater whereas a <sup>13</sup>C/<sup>12</sup>C enrichment factor of  $-23.0 \pm 2.1\%$  was obtained for degradation with site

34 groundwater alone. It can be concluded that biological degradation of both CS<sub>2</sub> contaminated soil and  
35 groundwater is likely to occur in the field suggesting that natural attenuation may be an appropriate remedial  
36 tool at some sites. The presence of biodegradation by-products including COS and H<sub>2</sub>S indicates that  
37 biodegradation of CS<sub>2</sub> is occurring and stable carbon isotopes are a promising tool to quantify CS<sub>2</sub> degradation.

38

## 39 **KEY WORDS**

40 *Carbon disulphide, carbon disulfide, biodegradation, microcosms, natural attenuation, stable*  
41 *carbon isotopes*

42

## 43 INTRODUCTION

44 Carbon disulphide (CS<sub>2</sub>) is a toxic, dense non-aqueous phase liquid (DNAPL) that is both  
45 highly volatile and highly flammable (Kalin et al., 2005). It is present in the environment due  
46 to anaerobic activity in sediments (Moret et al., 2000 and Lovelock, 1974), metabolism of  
47 naturally occurring sulphur compounds by soil bacteria and vegetation (Crookes et al., 1993),  
48 volcanic eruptions (Rasmussen et al., 1982), and the in-situ burning of hydrocarbon  
49 contaminated salt marsh (Devai et al., 1998). However, anthropogenic sources provide the  
50 primary source of CS<sub>2</sub> in the environment (Watts, 2000). Due to its high volatility and that it  
51 can ignite or explode when exposed to air (Kalin et al., 2005) remediation of CS<sub>2</sub>  
52 contaminated sites is difficult and hazardous. Therefore, the development of a remediation  
53 approach that removes CS<sub>2</sub> contamination from soil and groundwater without exposure to air  
54 is desirable.

55

56 Carbon disulphide has been produced commercially since 1880, and was used historically in  
57 a variety of industries including the viscose process (Beauchamp Jr. et al., 1983). In 1973  
58 approximately 65 million kilograms of CS<sub>2</sub> were released to the air in the US, whilst 35  
59 million kilograms reached water and land (SRI, 1975 cited in Peyton et al., 1976). Although  
60 demand for CS<sub>2</sub> has declined in recent years, it is predicted that the expanding viscose  
61 industries in Asia will increase CS<sub>2</sub> demand by approximately 4.7% in the period 2007 to  
62 2012 (Rojo et al., 2010). Carbon disulphide is also an intermediate formed during the  
63 degradation of carbon tetrachloride (CCl<sub>4</sub>) in granular sludge (van Eekert et al., 1998), and in  
64 a sandy aquifer under sulphate-reducing conditions (Devlin and Müller, 1999). Davis et al.  
65 (2003) reported CS<sub>2</sub> concentrations of up to 160 mg L<sup>-1</sup> on a CCl<sub>4</sub> contaminated site under  
66 highly reducing conditions due to abiotic degradation of CCl<sub>4</sub>. Given the above, it is

67 unsurprising that sites contaminated with CS<sub>2</sub> have been identified worldwide. In 2006, of  
68 the 1244 sites listed on the USEPA's National Priorities List (NPL), 139 sites had recorded  
69 CS<sub>2</sub> as a contaminant of concern (USEPA, 2006). This is a similar figure to the number of  
70 sites that have recorded the presence of other chlorinated solvents, such as CCl<sub>4</sub> (USEPA,  
71 2006).

72

73 A number of abiotic techniques for the in-situ remediation of CS<sub>2</sub> using zero-valent iron for  
74 groundwater (Kalin et al. 2005) and chemical oxidation for soil (Dulsey et al. 2001 and Ross  
75 et al. 2008) are available. However, to the authors' knowledge no investigations into natural  
76 attenuation of CS<sub>2</sub>, for contaminated land cleanup have been carried out. In order to  
77 demonstrate natural attenuation at a contaminated site, Monitored Natural Attenuation  
78 (MNA) protocols recommend a detailed site characterisation and assessment employing a  
79 'lines of evidence approach' (Morgan and Sinke, 2005). Primary evidence includes the  
80 demonstration that the contaminant plume is stable, shrinking or exhausted using historical  
81 contaminant concentrations. However, these data alone will not indicate whether a  
82 destructive attenuation mechanism is responsible for the decrease in concentrations (Carey et  
83 al., 2000). Geochemical and chemical data are often used as a secondary line of evidence to  
84 demonstrate whether a destructive process is causing attenuation. Secondary data includes  
85 the characterisation of known intermediates and products of biodegradation and compound  
86 specific isotope analysis (van Ras et al., 2007).

87

88 The degradation of CS<sub>2</sub> by microorganisms has been studied by a number of authors to  
89 investigate the potential for their use in waste gas treatment plants for manufacturing  
90 processes such as the viscose rayon process (Rothschild et al., 1969; Rajagopal and Daniels,  
91 1986; Ottengraf et al., 1986; Smith, 1988; Smith and Kelly, 1988; Kelly and Baker, 1990;

92 Kelly and Smith, 1990; Plas et al., 1993; Odintsova et al., 1993; Jordan et al., 1995; Jordan,  
93 1996; Jordan et al., 1997; Alcantara et al., 1999; Hartikainen et al., 2000; Sorokin et al., 2001;  
94 Pol et al., 2007). During the aerobic and anaerobic degradation of CS<sub>2</sub>, carbonyl sulphate  
95 (COS) and hydrogen sulphide (H<sub>2</sub>S) are both formed as intermediates (Smith and Kelly,  
96 1988; Jordan et al., 1997; Alcantara et al., 1999; Hartikainen et al., 2000; Sorokin et al., 2001;  
97 Pol et al., 2007). However, under anaerobic conditions COS is reported to accumulate to a  
98 greater extent, prior to being degraded (Smith and Kelly, 1988 and Pol et al., 1997). Smith  
99 and Kelly (1988) proposed that all the carbon in CS<sub>2</sub> is converted first to COS and then to  
100 CO<sub>2</sub> during both aerobic and anaerobic degradation (Equations 1 and 2).



103 Under aerobic conditions subsequent oxidation of H<sub>2</sub>S to elemental sulphur and eventually  
104 sulphate may also occur (Smith and Kelly, 1988 and Alcantara et al., 1999).

105

106 Compound specific stable isotope analysis measures the relative abundance of heavy and  
107 light isotopes in a compound (in this case <sup>12</sup>C and <sup>13</sup>C). Biological and abiotic reactions  
108 which break individual bonds tend to cause greater stable isotope fractionation than physical  
109 processes such as dilution, volatilisation and sorption which act on the whole molecule  
110 (Elsner et al., 2005). Therefore, stable isotope fractionation provides a powerful tool in  
111 determining whether the natural attenuation of xenobiotic compounds is occurring in the field  
112 (Sturchio et al., 1998; Hunkeler et al., 1999; Sherwood Lollar, 2001 and McKelvie et al.,  
113 2007). However, to the authors' knowledge, no studies have reported stable carbon isotope  
114 enrichment factors for the degradation of CS<sub>2</sub>. This study investigates the biologically  
115 mediated processes of natural attenuation of CS<sub>2</sub>, to provide information about CS<sub>2</sub>

116 degradation rates, identify whether COS and H<sub>2</sub>S are degradation intermediates and measure  
117 the carbon isotope signatures of degradation in CS<sub>2</sub> contaminated soils and groundwater.  
118 This information will assist determination of whether natural attenuation is occurring at CS<sub>2</sub>  
119 contaminated sites.

120

## 121 **MATERIALS AND METHODS**

### 122 **Chemicals and materials**

123 Experiments were carried out using general purpose reagent grade CS<sub>2</sub> (99.99% w/v, Hopkin  
124 and Williams).

125

### 126 **Soils and groundwater**

127 Soils were collected during remediation works at a former chemical manufacturing works in  
128 Stretford, Manchester, UK. Groundwater was collected from a former viscose rayon plant in  
129 Carrickfergus, Northern Ireland. All samples were transported and stored in the dark at 5°C  
130 until use. Further details of both sites are provided in Section 1 of the online resources. Soil  
131 samples contaminated with CS<sub>2</sub> DNAPL were exposed to a nitrogen atmosphere within a  
132 sealed glove bag to allow volatilisation of background CS<sub>2</sub> contamination. Soils were sieved  
133 to remove stones greater than 2.36 mm, prior to placing in thin walled plastic bags and  
134 storing in the anaerobic chamber at room temperature until ready for use.

135

136 Site groundwater was collected anaerobically in 1.92 L nitrogen purged glass jars from an  
137 area of known CS<sub>2</sub> contamination. Prior to sampling the borehole was purged until water  
138 quality parameters reached stable values. Groundwater was transferred under nitrogen to

139 collapsible Tedlar bags to ensure no headspace during preparation. The CS<sub>2</sub> concentration in  
140 the site groundwater, used in the unspiked tests, was approximately 87 mg L<sup>-1</sup>. Because  
141 higher initial concentrations were required to facilitate carbon isotope analysis of degradation  
142 products, site groundwater was spiked with a CS<sub>2</sub> stock solution prepared in methanol.  
143 Concentrations in the Tedlar bag for the spiked experiments were 250 mg L<sup>-1</sup> CS<sub>2</sub> and 157  
144 mg L<sup>-1</sup> methanol. Previous investigations had shown increased CS<sub>2</sub> degradation when  
145 sulphate was present (Cox et al., 2005), therefore Na<sub>2</sub>SO<sub>4</sub> (20 mM) was added in both tests.  
146 Full details of soil and groundwater preparation are included in Section 2 of the online  
147 resources.

148

#### 149 **Spiked and unspiked microcosm studies**

150 Microcosm studies were carried out in pre-sterilised 22 mL Chromacol glass vials. Soil (5 g  
151 ± 0.01 g) were added to each vial, along with 10 mL of site groundwater taken directly from  
152 the Tedlar bag using a Teflon and glass, gas tight syringe. Approximately 10.1 mL of  
153 headspace was present in each vial. Vials were sealed immediately with a Teflon faced  
154 aluminium crimp seal. Due to equilibration with the headspace in the vial, CS<sub>2</sub>  
155 concentrations in the water in the vials (C<sub>0</sub>) were 40 mg L<sup>-1</sup> (unspiked test) and 100 mg L<sup>-1</sup>  
156 (spiked test). All setup was undertaken in an anaerobic chamber (10% v/v H<sub>2</sub>, 5% v/v CO<sub>2</sub>  
157 and 85% N<sub>2</sub>).

158

159 Carbon disulphide free controls containing soil and groundwater (that had been exposed to a  
160 nitrogen atmosphere in a glove bag to remove CS<sub>2</sub> but spiked with methanol) were used to  
161 account for the microbial growth on methanol or any background carbon sources present.  
162 Groundwater microcosms containing CS<sub>2</sub> were set up to determine CS<sub>2</sub> losses due to abiotic  
163 and biological degradation within groundwater exclusively. Microcosms containing soil and



164 groundwater, referred to as soil microcosms from this point forward, were set up to  
165 investigate what additional losses could be attributed to the presence of microorganisms in  
166 the soils. Sterilisation of soil and groundwater using mercuric chloride ( $\text{HgCl}_2$ ) (final soil  
167 concentration of  $92 \text{ mg of Hg L}^{-1}$ ) was unsuccessful (Cox, 2008). Autoclaving for 90  
168 minutes on two occasions on consecutive days was required to successfully sterilise soil  
169 containing  $\text{CS}_2$  degrading bacteria. For details of sterilisation trials see Section 4 of the  
170 online resources. Summary details of the composition of controls and microcosms for both  
171 the unspiked and spiked tests are shown in Tables 5.1 and 5.2 in the online resources.

172

173 Soil microcosms were prepared in triplicate, while controls and microcosms containing  
174 groundwater only were prepared in duplicate. All vials were sealed with teflon faced  
175 aluminium crimp seals, wrapped in parafilm and stored in the dark in the anaerobic chamber  
176 at room temperature. Vials were removed from the anaerobic chamber at regular intervals  
177 and sampled sacrificially to minimise potential for losses due to volatilisation over 10 days.

178

## 179 **Analytical methods**

180 Analysis for  $\text{CS}_2$ ,  $\text{COS}$ ,  $\text{H}_2\text{S}$ ,  $\text{CO}_2$  and  $\text{CH}_4$  concentration was undertaken by GC-MS (Trace  
181 DSQ, Thermo Finnigan). Compound specific carbon isotope ratios ( $\delta^{13}\text{C}$ ) of  $\text{CS}_2$ ,  $\text{COS}$ ,  $\text{CO}_2$   
182 and  $\text{CH}_4$  in the vial headspace were measured using GC-C-IRMS (Isoprime, GV Instruments)  
183 for the spiked test only. All isotopes were reported using the delta notation referenced to  
184 Vienna Peedee Belemnite, VPDB. Detailed methods for all analyses are described in Section  
185 6 of the online resources.

186

187

188 **Quantification of isotope fractionation**

189 Fractionation is often quantified for comparison purposes using the Rayleigh relationship  
190 (Equation 3) (Mariotti et al., 1981).

191 
$$R = R_0 f^{(\alpha-1)} \quad (3)$$

192

193 where R is the isotopic ratio of the substrate,  $R_0$  is the initial isotopic ratio of the substrate, f  
194 is the remaining fraction of the substrate and  $\alpha$  is the fractionation factor. Equation 3 can be  
195 rearranged and expressed in  $\delta$ ‰ notation as shown in Equation 4 (Mariotti et al., 1981):

196

197 
$$\ln \left( \frac{\frac{\delta}{1000} + 1}{\frac{\delta_0}{1000} + 1} \right) = (\alpha - 1) \ln f = \frac{\varepsilon}{1000} \ln f \quad (4)$$

198

199 where  $\delta_0$  is the initial  $\delta$  value, and  $\varepsilon$  is the per mil enrichment factor, which represents the  
200 isotopic difference between the contaminant and its initial degradation product (Clark and  
201 Fritz, 1997). Equation 4 can be simplified to Equation 5 for small values of  $\delta$  (Mariotti et al.,  
202 1981).

203

204 
$$\delta - \delta_0 = \Delta\delta \cong 10^3 (\alpha - 1) \ln f = \varepsilon \ln f \quad (5)$$

205

206 and therefore a plot of change in  $\delta^{13}\text{C}$  against  $\ln(f)$  will be a straight line of gradient  $\varepsilon$  that  
207 goes through the origin.

208

## 209 RESULTS AND DISCUSSION

### 210 Degradation rates

211 The natural log of CS<sub>2</sub> concentrations (normalised with respect to initial concentration, C<sub>0</sub>)  
212 against time, are plotted in Figure 1 for soil microcosms (unspiked and spiked tests),  
213 sterilised soil with groundwater controls (spiked test) and microcosms containing  
214 groundwater-only (unspiked and spiked tests). Initial losses were considered by excluding  
215 the initial (time zero) data point and calculating the best fit line without specifying a y-  
216 intercept. Where consecutive sampling occasions showed that CS<sub>2</sub> concentrations were less  
217 than the limit of quantification (<0.008% v/v), the dataset has been modified to exclude the  
218 later sampling occasion, as including this data point skewed the linear regression.

219

220 In both the unspiked and spiked groundwater-only tests (Figure 1) carbon disulphide  
221 concentrations decreased by approximately 40%. Significantly more degradation was  
222 observed in soil microcosms where almost 100% degradation of CS<sub>2</sub> was observed in both  
223 the unspiked and spiked tests. Therefore the majority of CS<sub>2</sub> degradation was attributed to  
224 the biological activity within the soil. First-order degradation rate constants for soil  
225 microcosms in the unspiked test were calculated based on the modified datasets, as shown in  
226 Figure 1 and summarised in Table 1. In accordance with Equation 6, the rate constants for  
227 the unspiked and spiked soil microcosms ( $k_{\text{micro soil}}$ ) were corrected for CS<sub>2</sub> losses due to  
228 volatilisation, abiotic reactions and biodegradation from groundwater using the degradation  
229 rate constant for the spiked sterilised soil with groundwater controls ( $k_{\text{sterilised}}$ ). This gives a  
230 rate constant for the biodegradation due to soil microbes ( $k_{\text{degrad}}$ ) of  $>2.39 \pm 0.16 \times 10^{-2} \text{ h}^{-1}$  for  
231 the unspiked test and  $1.25 \pm 0.15 \times 10^{-2} \text{ h}^{-1}$  for the spiked test (Table 1).

232

$$k_{\text{degrad}} = k_{\text{microsoil}} - k_{\text{sterilised}} \quad (6)$$

233

234  $k_{\text{degrad}}$  for the spiked test is less than  $k_{\text{degrad}}$  from the unspiked test suggesting that the rate of  
235 degradation decreases with increasing initial  $\text{CS}_2$  concentration ( $C_0$  was  $40 \text{ mg L}^{-1}$  in the  
236 unspiked test and  $100 \text{ mg L}^{-1}$  in the spiked test) due to microbial inhibition. Similar  
237 inhibitory effects were recorded by Plas et al. (1993) at  $\text{CS}_2$  concentrations above  $150 \text{ mg L}^{-1}$ ,  
238 for degradation of  $\text{CS}_2$  by *Thiobacillus* K4, while Pol et al. (2007) found that  $\text{CS}_2$   
239 concentrations greater than  $22.8 \text{ mg L}^{-1}$  inhibited growth of *Thiomonas* sp. WZW.

240

#### 241 **Degradation products**

242 Carbonyl sulphide was not observed above the limit of quantitation (0.008% v/v) in any  
243 control vials or microcosms containing groundwater exclusively. However, following 30  
244 hours incubation COS was detected in both the  $\text{CS}_2$  unspiked and spiked soil microcosms at  
245 0.043 %v/v and 0.287 %v/v, respectively (Figure 2(a)). After 150 hours incubation the  
246 concentrations of COS in the spiked and unspiked soil microcosms were below the limit of  
247 quantification. If the biological degradation of 1 mole of  $\text{CS}_2$  generates 1 mole of COS  
248 (Equation 1), it would be expected that a 60% reduction in COS production would be  
249 observed correlating to the 60% reduction in the initial  $\text{CS}_2$  concentration. However, the  
250 observed reduction was 85%, and such discrepancies were attributed to the fact that the  
251 maximum COS concentrations may not have been recorded due to the 24-hour sampling  
252 interval.

253

254 As with the intermediate COS,  $\text{H}_2\text{S}$  was not detected above the limit of quantification  
255 (0.008% v/v) in control vials or microcosms containing groundwater exclusively. Hydrogen  
256 sulphide was also below the limit of quantification (0.008% v/v) in the unspiked soil  
257 microcosms (Figure 2(b)). However, in  $\text{CS}_2$  spiked microcosms containing soil,  $\text{H}_2\text{S}$

258 concentrations increased during the first 30 hours of the experiment, before decreasing to less  
259 than the limit of quantification at approximately 50 hours, and increasing again slightly at  
260 200 hours (Figure 2(b)). Therefore H<sub>2</sub>S is formed as an intermediate during the anaerobic  
261 degradation of CS<sub>2</sub> by soil microorganisms. Indeed, the second smaller peak observed in  
262 Figure 2(b) may indicate that H<sub>2</sub>S is also produced as a result of the subsequent degradation  
263 of COS, as proposed by Equation 2.

264

265 The generation of both COS and H<sub>2</sub>S as intermediates during the biodegradation of CS<sub>2</sub> is in  
266 accordance with the mechanism for CS<sub>2</sub> degradation proposed by Smith and Kelly (1988)  
267 (Equations 1 and 2). In tests under anaerobic conditions, they found that both COS and H<sub>2</sub>S  
268 accumulated (Smith and Kelly, 1988). Similarly, Pol et al., (2007) found that under  
269 anaerobic conditions, degradation of CS<sub>2</sub> by *Thiomonas* sp. WZW resulted in the  
270 accumulation of COS and H<sub>2</sub>S, which finally resulted in the inhibition of CS<sub>2</sub> degradation. It  
271 is unclear from our results whether the microorganisms responsible for the anaerobic  
272 degradation of CS<sub>2</sub> were responsible for the subsequent anaerobic degradation of COS and  
273 H<sub>2</sub>S (another microorganism in the mixed consortium may have caused this degradation).  
274 However it is encouraging that even under strictly anaerobic conditions COS and H<sub>2</sub>S did not  
275 accumulate to sufficient concentrations to significantly inhibit CS<sub>2</sub> degradation.

276

## 277 **Carbon isotope signatures**

278 Rayleigh plots for CS<sub>2</sub> carbon isotopes were constructed for all experiments spiked with CS<sub>2</sub>  
279 (Figure 3). Apart from two anomalous data points (circled in Figure 3(a)), most data points  
280 from the sterilised soil with groundwater controls are clustered around the x-axis (zero),  
281 revealing that significant fractionation is not occurring in these vials. Therefore, CS<sub>2</sub> losses  
282 in the sterilised soil control vials are mostly due to non-fractionating processes such as

283 volatilisation and sorption to soil and vials. The fractionation observed in the two anomalous  
284 data points coincides with a drop in CS<sub>2</sub> concentration and a slight increase in COS  
285 concentrations, indicating that CS<sub>2</sub> degradation may be occurring in these vials, possibly due  
286 to the presence of site groundwater or incomplete sterilisation of the soil.

287

288 The fractionation of carbon isotopes observed in microcosms containing groundwater  
289 exclusively (Figure 3(b)) and with soil (Figure 3(c)) follows a Rayleigh-type relationship.  
290 However, the carbon isotope enrichment factor for CS<sub>2</sub> degradation was different when soil  
291 was present. Initial losses were again considered by excluding the initial (time zero) data  
292 point and calculating the best fit line without specifying a y-intercept. An enrichment factor  
293 of  $-7.5 \pm 0.8\text{‰}$  was obtained for the soil microcosms (which contained both soil and  
294 groundwater), while an enrichment factor of  $-23.0 \pm 2.1\text{‰}$  was obtained for the less rapid, but  
295 highly fractionating degradation observed in microcosms with groundwater only.

296

297 Both enrichment factors obtained for CS<sub>2</sub> degradation are within the range of enrichment  
298 factors reported in literature for other organic compounds ( $-0.5\text{‰}$  to  $-32.1\text{‰}$ ) (Hunkeler et  
299 al., 2001a; Meckenstock et al., 2004; Sherwood Lollar et al., 1999; Ahad et al., 2000; Dayan  
300 et al., 1999; Hunkeler et al., 1999; Barth et al., 2002; Hunkeler et al., 2002; Hunkeler et al.,  
301 2001b). Of the many compounds studied previously, it would be expected that fractionation  
302 would be similar to that observed for short chain chlorinated hydrocarbons, as (1) compounds  
303 with greater numbers of carbon atoms would have “diluted” enrichment factors  
304 (Meckenstock et al., 2004), and (2) degradation in these compounds occurs by breaking the  
305 C-Cl bond, and chlorine is similar in atomic weight to sulphur (Elsner et al., 2005). Reported  
306 enrichment factors for biodegradation of chlorinated ethenes range from  $-7.1\text{‰}$  to  $-31.1\text{‰}$

307 (Dayan et al., 1999; Hunkeler et al., 1999; Barth et al., 2002; Hunkeler et al., 2002), which  
308 encompasses the enrichment factors found for degradation of CS<sub>2</sub>.

309

310 The Streitweiser Limit for breaking a C-S bond is 1.050 (Huskey, 1991). This is a  
311 semi-quantitative estimate of the maximum kinetic isotope effect (KIE) ( $1/\alpha$ ) that would be  
312 observed for a reaction that breaks a C-S bond (Elsner et al., 2005). This assumes bond  
313 cleavage at an infinitely late transition state, and therefore a more realistic estimate of the  
314 KIE may be obtained by assuming a transition state at 50% bond cleavage (Elsner et al.,  
315 2005), which corresponds to an estimated KIE of half the Streitweiser Limit (KIE = 1.025)  
316 (Elsner et al., 2005). KIEs and equivalent fractionation and enrichment factors estimated for  
317 C-S bond breakage and observed from CS<sub>2</sub> degradation are shown in Table 2.

318

319 Fractionation observed due to degradation with site groundwater only ( $\epsilon = -23.0 \pm 2.1\%$ )  
320 correlates well with the estimated values. The p-value for the gradient of the straight line  
321 regression on the Rayleigh plot for these vials is  $p = 1.4 \times 10^{-7}$ , indicating the null hypothesis  
322 that the straight line's true gradient is zero can be rejected comfortably (threshold p-value =  
323 0.05). Therefore the assumption that a linear relationship exists is acceptable for these  
324 results. The coefficient of determination,  $R^2$ , was greater than 0.90 (n=14), which is  
325 considered to be a good fit, given that vials were sacrificially sampled. A linear fit indicates  
326 that fractionation is controlled by a single reaction step (Ahad et al., 2000). The coefficient  
327 of determination,  $R^2$ , was also greater than 0.85 (n=17), for microcosms with soil, while the  
328 p-value for the gradient was  $p = 6.6 \times 10^{-8}$  ( $\epsilon = -7.5 \pm 0.8\%$ ), however the lower enrichment  
329 factor and higher reaction rate for these vials suggests the majority of degradation is  
330 occurring via a different pathway/mechanism than degradation due to site groundwater only.

331 This difference in fractionation factors is seemingly anomalous, and requires further  
332 investigation to fully elucidate.

333

334 Carbonyl sulphide was initially highly depleted in  $^{13}\text{C}$  in soil microcosms (as shown in Figure  
335 4), as it was being formed predominantly from  $\text{CS}_2$  molecules that contained  $^{12}\text{C}$  rather than  
336  $^{13}\text{C}$ . However as COS was subsequently degraded, its carbon isotope ratio became enriched,  
337 as the  $\text{CS}_2$  became enriched in  $^{13}\text{C}$  and concurrently COS molecules containing  $^{12}\text{C}$  were  
338 preferentially degraded. A similar trend in carbon isotope ratio was previously reported for  
339 intermediates produced during the degradation of chlorinated solvents (Hunkeler et al., 1999;  
340 Hunkeler et al., 2002).

341

#### 342 **Modelling of degradation product concentrations and isotope ratios**

343 To investigate the end point of degradation in soil microcosms, concentrations and isotope  
344 ratios of  $\text{CS}_2$  and potential degradation products ( $\text{COS}$ ,  $\text{CO}_2$  and  $\text{CH}_4$ ) from soil microcosms  
345 were modelled mathematically, using a method described by Hunkeler et al. (2002).  
346 Microbial and abiotic activity with groundwater only vials were not modelled, as COS  
347 concentrations in these vials were less than the limit of quantitation of the concentration  
348 analysis.

349

350 Two illustrative models were constructed, the first assuming that  $\text{CO}_2$  is the end point of  $\text{CS}_2$   
351 degradation, (Model 1, shown in Equation 7), and the second assuming that  $\text{CO}_2$  is removed  
352 from the system. In this case, it has been assumed that  $\text{CO}_2$  is converted to  $\text{CH}_4$  by  
353 methanogenesis (Model 2, shown in Equation 8). Both models treat the microcosms as  
354 closed systems, with no other carbon sources contributing to the production of COS,  $\text{CO}_2$  or  
355  $\text{CH}_4$ .





380 calculated by Models 1 and 2 shows that both models fit the CO<sub>2</sub> data within the error  
381 observed (Figure 8.2 in the online resources). CH<sub>4</sub> concentrations were also modelled  
382 adequately by Model 2 (Figure 8.3 of the online resources).

383

384 As Model 1 and Model 2 both fit the concentration data (see Figures 8.1 to 8.3 of the online  
385 resources), it is not possible from concentration data alone to determine whether the CH<sub>4</sub>  
386 produced in these vials was generated by methanogenic degradation of CO<sub>2</sub> or if another  
387 carbon source was degraded to form CH<sub>4</sub>. To investigate this further both models were  
388 extended to consider carbon isotope effects (see Section 7 of the online resources and Cox  
389 (2008) for full details).

390

391 Again the expressions for carbon isotope ratios for CS<sub>2</sub> and COS are the same for both Model  
392 1 and 2, and isotope ratios predicted by both models are shown on Figure 4. The small dip in  
393 CS<sub>2</sub> carbon isotope ratio observed at 100 hours may be due to an inhibitory affect caused by  
394 transitory accumulation of COS and H<sub>2</sub>S as Pol et al. (2007) found these intermediates can  
395 inhibit CS<sub>2</sub> degradation. The fit for COS does not appear to be as good as for CS<sub>2</sub>, however  
396 if any lag period was experienced before COS degradation commenced, this may explain the  
397 initial rise and fall, as the model would show more rapid enrichment over the first 100 hours,  
398 if the COS degradation rate was reduced.

399

400 Isotope data for CO<sub>2</sub> showed an enrichment in CO<sub>2</sub> isotope ratios of approximately 4‰ over  
401 the course of the experiment. This enrichment was not modelled by Model 1, but Model 2  
402 was able to replicate this (Figure 5(a)) if the CO<sub>2</sub> was degraded by a highly fractionating  
403 process, such as methanogenesis, which has a reported enrichment factor ( $\epsilon$ ) of  $-75 \pm 15\%$   
404 (Clarke and Fritz, 1997)). But modelling of CH<sub>4</sub> isotope data (Figure 5(b)) demonstrated that

405 this process was not methanogenic conversion of CO<sub>2</sub> to CH<sub>4</sub>. The actual CH<sub>4</sub> produced was  
406 initially depleted in <sup>13</sup>C, and rapidly became more enriched in <sup>13</sup>C. However, due to the low  
407 CO<sub>2</sub> degradation rate and high initial CO<sub>2</sub> concentration, the model predicts very slow  
408 enrichment if CO<sub>2</sub> was being converted to CH<sub>4</sub>. Therefore, as a rapid degradation rate would  
409 not fit the CH<sub>4</sub> concentration data, the CH<sub>4</sub> present in the vials must be produced as a result  
410 of degradation of another carbon source in the microcosm, such as methanol. This is  
411 supported by the fact that CH<sub>4</sub> was produced in CS<sub>2</sub> free controls in the spiked test (which  
412 contained methanol) but not CS<sub>2</sub> free controls in the unspiked tests (which did not contain  
413 methanol).

414

415 Therefore it is likely that CO<sub>2</sub> in the vials is being consumed, possibly by assimilation into  
416 the biomass of cells. Miltner et al. (2005) have suggested CO<sub>2</sub> fixation is a significant factor  
417 of microbial activity in soils. This could mean that the bacteria responsible for CS<sub>2</sub>  
418 degradation are obtaining energy from CS<sub>2</sub> and carbon from CO<sub>2</sub>, as described by Odintsova  
419 et al. (1993). However it is also possible that another microorganism is consuming CO<sub>2</sub>.

420

## 421 **CONCLUSIONS AND RECOMMENDATIONS**

422 Degradation experiments demonstrated that the soil tested contained indigenous bacteria that  
423 were capable of degrading CS<sub>2</sub>. This implies that natural attenuation could potentially be  
424 used to remediate CS<sub>2</sub> contaminated sites; however, further work is needed to characterise the  
425 conditions under which degradation is likely to occur in the field. Comparison of the rate  
426 constants calculated in both tests revealed that the first-order degradation rate constant  
427 decreases with increasing initial CS<sub>2</sub> concentration. This may be as a result of CS<sub>2</sub> being  
428 inhibitory to the CS<sub>2</sub> degrading organisms. Therefore site investigations should determine

429 whether natural attenuation would be restricted to down gradient portions of a CS<sub>2</sub> plume and  
430 whether source zone remediation would significantly enhance the performance of natural  
431 attenuation.

432

433 COS and H<sub>2</sub>S were both shown to be intermediates of anaerobic biodegradation of CS<sub>2</sub> by the  
434 bacteria present in the soil; however, no COS or H<sub>2</sub>S greater than the limit of quantitation  
435 was observed in control vials or during degradation with site groundwater alone. Therefore  
436 the presence of COS or H<sub>2</sub>S in groundwater may be good indicators that biodegradation of  
437 CS<sub>2</sub> is occurring in the field; however, their absence is not indicative that biodegradation is  
438 not occurring. Even under strictly anaerobic conditions COS and H<sub>2</sub>S did not accumulate to  
439 sufficient concentrations long enough to inhibit CS<sub>2</sub> degradation, which suggests that  
440 accumulation of by-products will not prevent natural attenuation from occurring in the field.

441

442 A <sup>13</sup>C/<sup>12</sup>C enrichment factor of  $-7.5 \pm 0.8\%$  was obtained for CS<sub>2</sub> degradation with both soil  
443 and site groundwater, whereas a <sup>13</sup>C/<sup>12</sup>C enrichment factor of  $-23.0 \pm 2.1\%$  was obtained for  
444 the less rapid degradation due to site groundwater alone, suggesting that if isotopic  
445 fractionation is observed in the field, it could indicate that degradation is occurring.  
446 However, as it appears that different mechanisms may cause differing amounts of  
447 fractionation, until a database of CS<sub>2</sub> enrichment factors has been established it will only be  
448 possible to quantify degradation once a site specific enrichment factor has been determined  
449 experimentally.

450

## 451 **ACKNOWLEDGEMENTS**

452 This work was supported by an industrial grant from AkzoNobel to Prof. Kalin. White Young Green  
453 Environmental is also acknowledged for obtaining permission to collect groundwater samples. The authors  
454 acknowledge the valuable technical assistance provided by Juliet McShane and Luc Belaid. We also thank Dr  
455 Debra Phillips and Dr Trevor Elliot for valuable advice and comments during preparation of this manuscript.

456

## 457 **REFERENCES**

- 458 Ahad JME, Sherwood Lollar B, Edwards EA, Slater GF, Sleep BE (2000) Carbon isotope fractionation during  
459 anaerobic biodegradation of toluene: Implications for intrinsic bioremediation. *Environ Sci Technol*  
460 34:892-896
- 461 Alcantara S, Estrada I, Vasquez MS, Revah S (1999) Carbon disulfide oxidation by a microbial consortium from  
462 a trickling filter. *Biotechnol Lett* 21:815-819
- 463 Ashworth J, Briggs GG, Evans AA, Matula J (1977) Inhibition of nitrification by nitrapyrin, carbon disulphide  
464 and trithiocarbonate. *J Sci Food and Agric* 28:673-683
- 465 Barth JAC, Slater GF, Schuth C, Bill M, Downey A, Larkin MJ, Kalin RM (2002) Carbon isotope fractionation  
466 during aerobic biodegradation of trichloroethene by *Burkholderia cepacia* G4: a tool to map  
467 degradation mechanisms. *Appl Environ Microbiol* 68(4):1728-1734
- 468 Beauchamp Jr. RO, Bus JS, Popp JA, Boreiko CJ, Goldberg L (1983) A critical review of the literature on  
469 carbon disulfide toxicity. *CRC Crit Rev Toxic* 11(3):169-278
- 470 Bond EJ (1984) *Manual of fumigation for insect control*. Research Centre Agriculture Canada, Ontario
- 471 Bremner JM, Banwart WL (1976) Sorption of sulfur gases by soils. *Soil Biol Biochem* 8(79):83
- 472 Carey MA, Finnamore JR, Morrey MJ, Marsland PA (2000) *Guidance on the assessment and monitoring of*  
473 *natural attenuation of contaminants in groundwater*. R&D Publication 95. Bristol, UK, Environment  
474 Agency
- 475 Cox SF, Kalin RM, McKinley JD, Morgan P, Thurgood R (2005) Preliminary investigations into biological  
476 processes of natural attenuation of carbon disulphide. Alleman BC and Kelley ME (eds). In:

477 Proceedings of the Eighth International In Situ and On-site Bioremediation Symposium. Battelle Press,  
478 Columbus, OH. Paper H3

479 Cox SF (2008) Investigation of the biological processes of natural attenuation of carbon disulphide in the  
480 ground. Dissertation, Queen's University Belfast

481 Clark I, Fritz P (1997) Environmental isotopes in hydrogeology. CRC Press Ltd, United States of America

482 Crookes MJ, Diment J, Dobson S (1993) Environmental hazard assessment: carbon disulphide. TSD14. Great  
483 Britain. Department of the Environment. Directorate of Air, Climate and Toxic Substances. Toxic  
484 Substances Division, BRE bookshop

485 Davis A, Fennemore GG, Peck C, Walker CR, McIlwraith J, Thomas S (2003) Degradation of carbon  
486 tetrachloride in a reducing groundwater environment: Implications for natural attenuation. Appl  
487 Geochem 18(4):503-525

488 Dayan H, Abrajano T, Sturchio NC, Winsor L (1999) Carbon isotope fractionation during reductive  
489 dehalogenation of chlorinated ethenes by metallic iron. Org Geochem 30:755-763

490 Devlin JF, Muller D (1999) Field and laboratory studies of carbon tetrachloride transformation in a sandy  
491 aquifer under sulfate reducing conditions. Environ Sci Technol 33:1021-1027

492 Devai I, DeLaune RD, Henry CB, Lindau CW, Roberts PO (1998) Environmental significance of atmospheric  
493 emission resulting in *in situ* burning of oiled salt marsh. Mar Environ Res 45(2):157-167

494 Dulsey R, Mausner M, Bybee G, Mercurio B, Fahrenthold P, Hazard J, McGhee C (2001) Method for oxidative  
495 destruction of carbon disulfide in soil (US 6283675 B1)

496 Elsner M, Zwank L, Hunkeler D, Schwarzenbach RP (2005) A new concept linking observable stable isotope  
497 fractionation to transformation pathways of organic pollutants. Environ Sci Technol 39:6896-6916

498 Gelbke H, Goen T, Maurer M, Sulsky SI (2009) A review of health effects of carbon disulphide in viscose  
499 industry and a proposal for an occupational exposure limit. Crit Rev Toxicol. 39(S2):1-126

500 Gustafsson JP (2006) Help for Visual Minteq 2.5, KTH, Department of Land and Water Resources Engineering.  
501 Available with Visual Minteq 2.5.

502 Gustafsson JP (2012) Visual Minteq, KTH, Department of Land and Water Resources Engineering.  
503 <http://www2.lwr.kth.se/English/OurSoftware/vminteq/>. (Accessed 22 June 2012)

504 Hartikainen T, Ruuskanen J, Raty K, von Wright A, Martikainen PJ (2000) Physiology and taxonomy of  
505 Thiobacillus strain TJ330, which oxidizes carbon disulphide (CS<sub>2</sub>). Journal of Appl Microbiol  
506 89(4):580-586

507 Hunkeler D, Andersen N, Aravena R, Bernasconi SM, Butler BJ (2001a) Hydrogen and carbon isotope  
508 fractionation during aerobic biodegradation of benzene. *Environ Sci Technol* 35:3462-3467

509 Hunkeler D, Aravena R, Butler BJ (1999) Monitoring microbial dechlorination of tetrachloroethene (PCE) in  
510 groundwater using compound-specific stable carbon isotope ratios: Microcosm and field studies.  
511 *Environ Sci Technol* 33:2733-2738

512 Hunkeler D, Aravena R, Cox E (2002) Carbon isotopes as a tool to evaluate the origin and fate of vinyl chloride:  
513 Laboratory experiments and modelling of isotope evolution. *Environ Sci Technol* 36:3378-3384

514 Hunkeler D, Butler BJ, Aravena R, Barker JF (2001b) Monitoring biodegradation of methyl tert-butyl ether  
515 (MTBE) using compound-specific carbon isotope analysis. *Environ Sci Technol* 35:676-681

516 Huskey WP (1991) Origins and interpretations of heavy-atom isotope effects. In: Cook PF (ed) *Enzyme  
517 mechanism from isotope effects*, 1st edn. CRC Press, Boca Raton, FL, pp 47-72

518 Jordan SL (1996) *Physiology and biochemistry of bacteria which oxidise carbon disulphide*. Dissertation,  
519 University of London

520 Jordan SL, Kraczkiewicz-Dowjat AJ, Kelly DP, Wood AP (1995) Novel eubacteria able to grow on carbon  
521 disulfide. *Arch Microbiol* 163:131-137

522 Jordan SL, McDonald IR, Kraczkiewicz-Dowjat AJ, Kelly DP, Rainey FA, Murrell JC, Wood AP (1997)  
523 Autotrophic growth on carbon disulfide is a property of novel strains of *Paracoccus denitrificans*. *Arch  
524 Microbiol* 168:225-236

525 Kalin RM, Myles P, McGeough KL (2005) Carbon disulphide sources, risks and treatment: An emerging  
526 contaminant of environmental concern? Boshoff GA and Bone BD (eds). In: *Permeable Reactive  
527 Barriers*. IAHS Press. 82-92

528 Kelly DP, Baker SC (1990) The organosulphur cycle: aerobic and anaerobic processes leading to turnover of C<sub>1</sub>-  
529 sulphur compounds. *FEMS Microbiol Rev* 87:241-246

530 Kelly DP, Smith NA (1990) Organic sulfur compounds in the environment. *Biogeochemistry, microbiology and  
531 ecological aspects*. *Adv Microb Ecol*. 11:345-385

532 Lovelock JE, (1974) CS<sub>2</sub> and the natural sulphur cycle. *Nat* 248:625-626

533 Ma J, Ji J, Ding Q, Liu W, Wang S, Wang N, Chem G (2010) The effects of carbon disulfide on male sexual  
534 function. *Toxicol Ind Health*. 26(6):375-382

535 Mariotti A, Germon JC, Hubert P, Kaiser P, Letolle R, Tardieux A, Tardieux P (1981) Experimental  
536 determination of nitrogen kinetic isotope fractionation: some principles; illustrations for the  
537 denitrification and nitrification processes. *Plant Soil* 62:413-430

538 McKelvie JR, Mackay DM, Sieyes NR, Lacrampe-Couloume G, Sherwood Lollar B (2007) Quantifying MTBE  
539 biodegradation in Vandenberg Air Force Base ethanol release study using stable carbon isotopes. *J*  
540 *Contam Hydrol* 94:157-165

541 Meckenstock RU, Morasch B, Griebler C, Richnow HH (2004) Stable isotope fractionation analysis as a tool to  
542 monitor biodegradation in contaminated aquifers. *J Contam Hydrol* 75:215-255

543 Miltner A, Kopinke F, Kindler R, Selesi D, Hartmann A, Kastner M (2005) Non-phototrophic CO<sub>2</sub> fixation by  
544 soil microorganisms. *Plant Soil* 269:193-20

545 Moret I, Gambaro A, Piazza R, Barbante C, Andreoli C, Corami F, Scarponi G (2000) The seasonal variations  
546 of dimethyl sulphide and carbon disulphide in surface waters of the Venice lagoon. *Mar Chem* 71(3-  
547 4):283-295

548 Morgan P, Sinke A (2005) Monitored natural attenuation. *Land Contam Reclam* 13(3):247-265

549 Odinstova EV, Wood AP, Kelly DP (1993) Chemolithoautotrophic growth of *Thiothrix ramosa*. *Arch Microbiol*  
550 160:152-157

551 Ottengraf SPP, Meesters JJP, van den Oever, A. H. C., Rozema HR (1986) Biological elimination of volatile  
552 xenobiotic compounds in biofilters. *Bioprocess Eng* 1:61-69

553 Peyton TO, Steele RV, Mabey WR (1976) Carbon Disulfide, Carbonyl Sulfide. Literature Review and  
554 Environmental Assessment. EPA-600/9-78-009. Washington DC, United States Environmental  
555 Protection Agency. Office of Health and Ecological Effects

556 Plas C, Wimmer K, Holubar P, Mattanovich D, Danner H, Jelinek E, Harant H, Braun R (1993) Degradation of  
557 carbon disulphide by a *Thiobacillus* isolate. *Appl Microbiol Biotechnol* 38(6):820-823

558 Pol A, van der Drift C, Op den Camp, H. J. M. (2007) Isolation of carbon disulfide utilizing *Thiomonas* sp. And  
559 its application in a biotrickling filter. *Appl Microb Cell Physiol* 74:439-446

560 Rajagopal BS, Daniels L (1986) Investigation of mercaptans, organic sulfides, and inorganic sulfur compounds  
561 as sulfur sources for the growth of methanogenic bacteria. *Curr Microbiol* 14:137-144

562 Rasmussen RA, Khalil MA, Dalluge RW (1982) Carbonyl sulfide and carbon disulfide from the eruptions of  
563 Mount St. Helens. *Sci* 215:665-667



564 Rojo N, Gallastegi G, Barona A, Gurtubay L, Ibarra-Berastegi G, Elias A (2010) Biotechnology as a alternative  
565 for carbon disulfide treatment in air pollution control. *Environ Rev* 18:321-332

566 Ross I, MacLeod C, Harvey B, Burdick J, Maggs D (2008) In-situ remediation of carbon disulphide. In:  
567 Consoil 2008 – CDROM edition: Proceedings of the 10<sup>th</sup> International UFZ-Deltares/UNO Conference  
568 on Soil-Water Systems. E474-E475

569 Rothschild B, Butler R, Keller JR (1969) Metabolism of carbon disulfide by *Thiobacillus thiooxidans*. *Bacteriol*  
570 *Proc*:64

571 Sherwood Lollar B, Slater GF, Ahad J, Sleep B, Spivack J, Brennan M, MacKenzie P (1999) Contrasting carbon  
572 isotope fractionation during biodegradation of trichloroethylene and toluene: implications for intrinsic  
573 bioremediation. *Org Geochem* 30:813-820

574 Sherwood Lollar B, Slater GF, Sleep B, Witt M, Klecka GM, Harkness M, Spivack J (2001) Stable carbon  
575 isotope evidence for intrinsic bioremediation of tetrachloroethene and trichloroethene at Area 6, Dover  
576 Air Force Base. *Environ Sci Technol* 35:261-269

577 Smith NA (1988) Metabolism of dimethyl sulphide, carbon disulphide and other volatile sulphur. Dissertation,  
578 University of Warwick

579 Smith NA, Kelly DP (1988) Oxidation of carbon disulphide as the sole source of energy for the autotrophic  
580 growth of *Thiobacillus thioparus* strain TK-m. *J Gen Microbiol* 134(11):3041-3048

581 Sorokin DY, Tourova TP, Lysenko AM, Kuenen JG (2001) Microbial Thiocyanate Utilisation under Highly  
582 Alkaline Conditions. *Appl Environ Microbiol* 67(2):528-538

583 Sturchio NC, Clausen JL, Heraty LJ, Huang L, Holt BD, Abrajano JR (1998) Chlorine isotope investigation of  
584 natural attenuation of trichloroethene in an aerobic aquifer. *Environ Sci Technol* 32(20):3037-3042

585 U.S. Environmental Protection Agency (USEPA) (2006) National Priorities List (NPL): Advanced query form.  
586 U.S. Environmental Protection Agency (USEPA).  
587 <http://www.epa.gov/superfund/sites/query/advquery.htm>. Accessed August, 4 2006

588 Van Eekert, M. H. A., Schroder TJ, Stams AJM, Schraa G, Field JA (1998) Degradation and fate of carbon  
589 tetrachloride in unadapted methanogenic granular sludge. *Appl Environ Microbiol* 64(7):2350-2356


590 van Ras, N. J. P., Winters RO, Lieten SH, Dijkhuis JE, Henssen MJC, van Hattem WA, Lethbridge G (2007)  
591 Sustainability of natural attenuation of aromatics (BTEX). 2003.1969. Rotterdam, Network for  
592 industrially contaminated land in Europe (NICOLE)

593 Watts SF, (2000) The mass budgets of carbonyl sulfide, dimethyl sulfide, carbon disulfide and hydrogen sulfide.  
594 Atmos Environ 34:761-779.

595 Wilson JT, Kaiser PM, Adair C (2005) Monitored Natural Attenuation of MTBE as a Risk Management Option  
596 at Leaking Underground Storage Tank Sites. EPA/600/R-04/1790. Cincinnati, OH, National Risk  
597 Management Research Laboratory, Office of Research and Development, USEPA  
598

599 **Figures**

600




601


602 **Fig. 1** Plot of  $\ln$  normalised  $\text{CS}_2$  concentration versus time for an initial  $\text{CS}_2$  concentration of  $40 \text{ mg L}^{-1}$   
 603 (unspiked test) and  $100 \text{ mg L}^{-1}$  (spiked test). Error bars are two standard errors of three replicate measurements,  
 604 and therefore depict error associated with method of analysis. LoQ is limit of quantification

605

606




607




608 **Fig. 2** Headspace concentration versus time for an initial CS<sub>2</sub> concentration of 40 mg L<sup>-1</sup> (unspiked test) and  
609 100 mg L<sup>-1</sup> (spiked test) for (a) COS and (b) H<sub>2</sub>S. Limit of Quantification (LoQ) is 0.008% v/v for both COS  
610 and H<sub>2</sub>S. Error bars are two standard errors of three independent samples


611



612




613



614 **Fig. 3** Rayleigh plot of  $\Delta\delta^{13}\text{C}$  versus  $-\ln(f)$  for (a) sterilised soil control, (b) groundwater microcosms (p-value  
615 (gradient) =  $1.4 \times 10^{-7}$ ) and (c) groundwater and soil microcosms (p-value (gradient) =  $6.6 \times 10^{-8}$ ). Circles in  
616 Fig.3(a) identify two anomalous data points

617

618




619

620 **Fig. 4** Actual and modelled carbon isotope ratio of CS<sub>2</sub> and COS versus time for microcosms with soil and  
621 site groundwater. Error bars are two standard errors of three independent samples (except t = 191 hrs and 240  
622 hrs, where n = 1 (for CS<sub>2</sub>) and t = 148 hrs where n = 2 (for COS))


623

624

625



626



627 **Fig 5** Actual and modelled carbon isotope data versus time for  $k_1 = 0.0163 \text{ h}^{-1}$ ,  $\alpha_1 = 0.9919$ ,  $k_2 = 0.15 \text{ h}^{-1}$ ,  
628  $\alpha_2 = 0.985$ ,  $k_3 = 0.00055 \text{ h}^{-1}$  and  $\alpha_3 = 0.925$  for (a)  $\text{CO}_2$  and (b)  $\text{CH}_4$ . Error bars are two standard errors of three  
629 independent samples

630

631

632 **Tables**

633

<b>Sterilised soil control (<math>C_0 = 100\text{mg L}^{-1}</math>)</b>	$k_{\text{sterilised}} (\text{h}^{-1})$	$0.15 \pm 0.04 \times 10^{-2}$
<b>Groundwater only (<math>C_0 = 40\text{mg L}^{-1}</math>)</b>	$k_{\text{groundwater}} (\text{h}^{-1})$	$0.26 \pm 0.12 \times 10^{-2}$
<b>Groundwater only (<math>C_0 = 100\text{mg L}^{-1}</math>)</b>	$k_{\text{groundwater}} (\text{h}^{-1})$	$0.38 \pm 0.11 \times 10^{-2}$
<b>Soil and groundwater (<math>C_0 = 40\text{mg L}^{-1}</math>)</b>	$k_{\text{micro soil}} (\text{h}^{-1})$	$>2.54 \pm 0.15 \times 10^{-2}$
<b>Soil and groundwater (<math>C_0 = 100\text{mg L}^{-1}</math>)</b>	$k_{\text{micro soil}} (\text{h}^{-1})$	$1.40 \pm 0.14 \times 10^{-2}$

634 **Table 1** First-order degradation rate constants ( $\text{h}^{-1}$ ) for unspiked and spiked tests. Uncertainties are one  
 635 standard error

636

	<b>KIE</b>	<b><math>\alpha</math></b>	<b><math>\epsilon</math> (‰)</b>
<b>Steitweiser limit for C-S bond<sup>a</sup></b>	1.050	0.952	-48
<b>More realistic estimate of KIE<sup>a</sup></b>	1.025	0.976	-24
<b>Degradation due to site groundwater</b>	1.0235	0.9770	-23.0
<b>Degradation due to soil and site groundwater</b>	1.0076	0.9925	-7.5

637 <sup>a</sup> taken from Huskey (1991)

638 **Table 2** KIE, fractionation factor ( $\alpha$ ) and enrichment factor ( $\epsilon$ ) calculated for C-S bond breakage and observed  
 639 during degradation of  $\text{CS}_2$