



**QUEEN'S
UNIVERSITY
BELFAST**

Comment on “Mathematical and physical aspects of Kappa velocity distribution” [Phys. Plasmas 14, 110702 (2007)]

Hellberg, M. A., Mace, R. L., Baluku, T. K., Kourakis, Y., & Saini, N. (2009). Comment on “Mathematical and physical aspects of Kappa velocity distribution” [Phys. Plasmas 14, 110702 (2007)]. *Physics of Plasmas*, 16(9), Article 094701. Advance online publication. <https://doi.org/10.1063/1.3213388>

Published in:
Physics of Plasmas

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights
© 2009 American Institute of Physics

General rights
Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Open Access
This research has been made openly available by Queen's academics and its Open Research team. We would love to hear how access to this research benefits you. – Share your feedback with us: <http://go.qub.ac.uk/oa-feedback>

Comment on “Mathematical and physical aspects of Kappa velocity distribution” [Phys. Plasmas 14, 110702 (2007)]

M. A. Hellberg,^{1,a)} R. L. Mace,^{1,b)} T. K. Baluku,^{1,c)} I. Kourakis,^{2,d)} and N. S. Saini^{2,e)}

¹School of Physics, University of KwaZulu-Natal, Private Bag X54001, Durban 4000, South Africa

²Department of Physics and Astronomy, Centre for Plasma Physics, Queen’s University Belfast, Belfast, Northern Ireland BT7 1NN, United Kingdom

(Received 27 April 2009; accepted 7 August 2009; published online 3 September 2009; publisher error corrected 10 September 2009)

A recent paper [L.-N. Hau and W.-Z. Fu, Phys. Plasmas 14, 110702 (2007)] deals with certain mathematical and physical properties of the kappa distribution. We comment on the authors’ use of a form of distribution function that is different from the “standard” form of the kappa distribution, and hence their results, *inter alia* for an expansion of the distribution function and for the associated number density in an electrostatic potential, do not fully reflect the dependence on κ that would be associated with the conventional kappa distribution. We note that their definition of the kappa distribution function is also different from a modified distribution based on the notion of nonextensive entropy. © 2009 American Institute of Physics. [doi:10.1063/1.3213388]

In a recent paper,¹ a number of mathematical aspects of the kappa velocity distribution² are presented. We are concerned that the results of Hau and Fu¹ do not correctly represent the full dependence on the parameter κ because the authors do not use the “conventional” kappa distribution² in their calculations.

The kappa distribution has been commonly used to fit particle data obtained in satellite-based experiments, for instance, as discussed by Refs. 2–8, as well as in some laboratory devices.⁹ It describes a distribution function that has a Maxwellian-like core and a high-energy component of power-law form. The conventional isotropic, three-dimensional form of the kappa distribution function may be written concisely as

$$F_{\kappa}(v) = A_{\kappa} \left(1 + \frac{v^2}{\kappa\theta^2} \right)^{-(\kappa+1)}. \quad (1)$$

This expression contains three parameters, viz., A_{κ} , θ , and κ , the latter being a spectral index.^{2,10,11} This is essentially the power index obeyed by the high-energy component of the equivalent energy distribution. Of these parameters, only κ is in a sense a free parameter, as the other two are constrained by the lowest (even) moments of the distribution function. Equating the lowest moment to the density N leads to

$$A_{\kappa} = \frac{N}{(\pi\kappa\theta^2)^{3/2}} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa-1/2)}, \quad (2)$$

where $\Gamma(x)$ is the usual gamma function. The parameter θ is a characteristic speed, that has been termed an effective thermal speed by some authors. In fact, given the isotropic distribution function in the form of Eq. (1), it can easily be shown that θ is the most probable particle speed. However,

we note that this is not the most probable speed that is found for a Maxwellian, i.e., what is usually termed the “thermal speed,” $v_t = (2K_B T/m)^{1/2}$, where T is the particle temperature, m the mass of the particles, and K_B is Boltzmann’s constant.

It is our contention that to properly relate θ to the plasma temperature requires consideration of the second moment of the velocity distribution function and some care in its interpretation.

To illustrate the source of the divergent approaches to the use of the kappa distribution, we provide here a brief historical perspective.

The kappa distribution was first used as a fit to particle data found with the OGO-1 satellite by Vasyliunas,² following a suggestion by Olbert.² In that paper, the standard isotropic three-dimensional distribution was written as²

$$F_{\kappa}(v) = \frac{N}{w_0^3} \frac{\Gamma(\kappa+1)}{(\pi\kappa)^{3/2} \Gamma(\kappa-1/2)} \left(1 + \frac{v^2}{\kappa w_0^2} \right)^{-(\kappa+1)}. \quad (3)$$

Here w_0 was stated to be the most probable particle speed, in agreement with our statement regarding the equivalent θ in Eq. (1).

Vasyliunas² also pointed out that it was often convenient to use the characteristic energy, $E_0 = mw_0^2/2$, corresponding to the characteristic speed w_0 . This is the energy of the peak in the differential flux. By considering the second moment of the distribution function,

$$U = \int dv f(v) m v^2 / 2, \quad (4)$$

it was shown that the mean energy per particle is related to the most probable speed w_0 through the associated energy E_0 by

$$E_m \equiv U/N = \frac{3}{2} E_0 \left(\frac{\kappa}{\kappa - 3/2} \right). \quad (5)$$

Finally, it was pointed out that in the special case $\kappa \rightarrow \infty$, the velocity distribution (3) reduces to the Maxwellian, for

^{a)}Electronic mail: hellberg@ukzn.ac.za.

^{b)}Electronic mail: macer@ukzn.ac.za.

^{c)}Electronic mail: 205527176@ukzn.ac.za and balukuthomas@yahoo.co.uk.

^{d)}Electronic mail: i.kourakis@qub.ac.uk.

^{e)}Electronic mail: nssaini@yahoo.com; ns.saini@qub.ac.uk.

which equipartition of energy applies rigorously, and the expression (5) reduces to $K_B T = E_0$.

A few years later, Formisano *et al.*³ used the kappa distribution (they called it a “ K distribution”) to fit proton data measured in the magnetosheath. Substituting the above notation for theirs, we note that they introduced a “plasma temperature T ,” which they expressed as

$$K_B T = (1/2) m v_0^2 \left(\frac{\kappa}{\kappa - 3/2} \right). \quad (6)$$

Clearly, this is related to the mean energy per particle, E_m , of Vasyliunas, by $E_m = (3/2) K_B T$. Such a temperature definition, making use of equipartition of energy, although appropriate for the equilibrium distribution, the Maxwellian, is not strictly valid for a non-Maxwellian distribution. Nevertheless, there are practical advantages to using such an equivalent kinetic temperature, which can be a useful concept, and moreover, is accepted in practice for distributions that do diverge from the Maxwellian.¹²

In what may be the earliest theoretical paper based on the kappa distribution, Leubner¹³ considered resonant wave particle interactions in a kappa plasma. Considering an anisotropic plasma ($T_\perp \neq T_\parallel$), he wrote the kappa distribution function in a form equivalent to that of Vasyliunas, introducing θ to represent the characteristic (most probable) speed. He also related the average energy to a temperature, hence obtaining

$$\theta^2 = \left(\frac{\kappa - 3/2}{\kappa} \right) \left(\frac{2K_B T}{m} \right). \quad (7)$$

Thus θ , in Leubner’s early paper, is explicitly not the Maxwellian thermal speed v_T . Furthermore, it is clearly κ -dependent. This is true also of the equivalent characteristic speed w_0 in Refs. 2 and 3.

Subsequently, a number of authors have studied various aspects and applications of the kappa distribution. Considering noise in a kappa plasma, Chateau and Meyer-Vernet¹⁴ calculated “the equivalent temperature T ” from the mean particle energy, obtaining the expression

$$T = \frac{\langle m v^2 \rangle}{3K_B} = \frac{m v_0^2}{K_B} \frac{\kappa}{2\kappa - 3}, \quad (8)$$

where they used v_0 in place of the notation w_0 used by Vasyliunas,² and the distribution is normalized by setting $\langle v^0 \rangle = 1$.

It is important to note that the use of equipartition of energy implies that the kinetic temperature (defined through the second moment of the distribution function) is the same for all plasmas with the same mean kinetic energy per particle, independently of the exact form of the distribution function. In our case, it is thus independent of the value of κ , as one can see from Eq. (8). The second equality of Eq. (8) stresses the relationship between the temperature T and the most probable speed (v_0) [i.e., θ in Eq. (7)] that is given explicitly in Eq. (7).

Independently, in a landmark set of papers, Summers, Thorne *et al.*^{10,11,15–19} considered various waves in such a kappa plasma. They introduced a modified plasma dispersion

function Z_κ^* . This is the equivalent, for waves in a kappa-distributed plasma, of the usual Z -function²⁰ for a Maxwellian distribution. In particular, they used the notation of Leubner,¹³ writing the isotropic, three-dimensional distribution function as¹⁰

$$F_\kappa(v) = \frac{N}{\pi^{3/2}} \frac{1}{\theta^3} \frac{\Gamma(\kappa + 1)}{\kappa^{3/2} \Gamma(\kappa - 1/2)} \left(1 + \frac{v^2}{\kappa \theta^2} \right)^{-(\kappa+1)}, \quad (9)$$

where $\theta = [(2\kappa - 3)/\kappa]^{1/2} (K_B T/m)^{1/2}$, with T the “particle temperature.” The latter was defined in terms of the mean particle kinetic energy, as shown explicitly, for instance, for anisotropic plasmas.^{16,18} We note that a well-defined value of θ requires $\kappa > 3/2$.^{10,21}

The formulation set out above has been followed by numerous papers on waves in kappa plasmas and related calculations, e.g., Refs. 7, 9, and 21–44 and many others.

From the above it also follows that one finds that the shielding distance in a κ -distributed plasma becomes $\lambda_\kappa = \lambda_D [(\kappa - 3/2)/(\kappa - 1/2)]^{1/2}$, where $\lambda_D = (\epsilon_0 K_B T / n e^2)^{1/2}$ is the usual Debye length for a Maxwellian plasma having the same density and the same mean energy.^{6,14,25}

As an aside, we note that integration over two velocity space coordinates leads to the one-dimensional kappa distribution function,^{10,27,38} which can be written as

$$f_\kappa(v) = \frac{N_0}{(\pi \kappa \theta^2)^{1/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa - 1/2)} \left(1 + \frac{v^2}{\kappa \theta^2} \right)^{-\kappa}, \quad (10)$$

where it is seen that the inverse power is κ , whereas for the three-dimensional distribution it is $(\kappa + 1)$.

Substituting in $F_\kappa(v)$ for θ , one may write the kappa distribution in an alternative form,³² i.e.,

$$F_\kappa(v) = \frac{N_0}{(\pi \kappa \theta^2)^{3/2}} \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa - 1/2)} \times \left(1 + \frac{v^2}{[\kappa - 3/2][2K_B T/m]} \right)^{-(\kappa+1)}. \quad (11)$$

In some instances the “characteristic velocity” (the most probable speed) is loosely referred to as a “thermal velocity.” However, in all the above cases, one demands of the distribution function that the parameters A_κ and θ should be obtained in a self-consistent manner from the lowest two even moments of the distribution, in which case it is clearly shown that, for a given value of mean particle energy, a measurable quantity, θ must be a function of κ . As made explicit in Ref. 6, one is actually comparing kappa distributions having different values of κ with a Maxwellian counterpart having the same number density and energy density—what one may call “the equivalent Maxwellian.” Although one may have qualms about applying equipartition to a nonequilibrium distribution, the interpretation of T as being the temperature of the equivalent Maxwellian is valid and is useful, and it is accepted in practice. Sometimes this T is called the kinetic temperature.

We note that, as the factor $(\kappa/[\kappa - 3/2]) > 1$ for finite κ , it follows from Eq. (8) that the “typical speed” v_0 of the conventional kappa distribution is smaller than the Maxwellian thermal speed for the same value of temperature. From

that and from Eq. (9) it follows that the standard kappa distribution has a higher, narrower peak, and a broader base than the equivalent Maxwellian. This is intuitively reasonable—if the total energy content is the same, and the kappa distribution has more high-energy particles, then the velocity spread of the core, and hence the *perceived* thermal core temperature, of the kappa distribution must be lower than that of the equivalent Maxwellian. Clearly, however, from the definition of temperature, all distributions with the same mean energy per particle have the same temperature, as we have seen above.

In our discussion below we prefer not to discuss a “temperature” of the kappa distribution as such, but instead shall concentrate on distributions having the same average energy density (the second moment of the distribution), a measurable quantity, and write the temperature of the equivalent Maxwellian. We shall then explore the effects of different values of κ on various characteristics of the plasma.

It obviously makes practical and theoretical sense to compare the kappa distribution to a Maxwellian distribution that has the same average kinetic energy per particle. As we have seen, that leads to the result, Eq. (7), that the characteristic velocity (the most probable speed θ) is proportional to the thermal speed of the equivalent Maxwellian. The constant of proportionality between θ and v_t is clearly dependent on κ . We reiterate this important point for future reference.

In contrast to the above formulation of the conventional kappa distribution, Leubner has, in a set of recent papers,^{45–47} followed on the work of Silva *et al.*⁴⁸ to suggest a modified kappa distribution, which is based on the notion of nonextensive entropy. It assumes a simple transformation from a Tsallis distribution.⁴⁹ The latter is expressed in terms of a parameter q , and writing $\kappa=1/(1-q)$, Leubner⁴⁵ obtained a one-dimensional distribution [Eq. (6) of Ref. 45]. In particular, in it, the most probable particle speed θ is replaced by the most probable speed of a Maxwellian, the thermal speed, $v_t=(2K_B T/m)^{1/2}$, where T is the temperature “of the species considered.” He then states that, by “analogy,” one finds a similar isotropic three-dimensional distribution [Eqs. (10) and (11) of Ref. 45],

$$F_L(\mathbf{v}) = \frac{N}{\pi^{3/2} v_t^3} \frac{1}{\kappa^{3/2}} \frac{\Gamma(\kappa)}{\Gamma(\kappa-3/2)} \left(1 + \frac{v^2}{\kappa v_t^2}\right)^{-\kappa}. \quad (12)$$

We note, importantly, that this three-dimensional function involves an inverse power κ , not $(\kappa+1)$. Later in the same paper, Leubner writes what he terms the “standard kappa distribution,” “written conventionally” in a form that is at first sight akin to that of Eq. (9), including an inverse power $(\kappa+1)$. However, unfortunately, he replaces θ by the thermal speed v_t . He later states that an “effective thermal speed $\theta=v_t[(\kappa-3/2)/\kappa]^{1/2}$ is commonly defined from the moments of the distribution function,” but does not use it. In a later paper,⁴⁶ similar results are put forward and lead to a distribution of the form of Eq. (12). In that paper, too, he writes of the “conventional” form of the kappa distribution function [alluding to that of Eq. (9)], but wrongly writes it in terms of the thermal speed v_t rather than the most probable speed θ . In Ref. 47, a paper cited in the Response⁵⁰ to our Comment,

essentially the same approach is taken. That is, an equivalent of Eq. (12) appears, and the misleading representation (use of v_t) of the conventional kappa distribution¹⁰ persists.

Although what is in essence a “ q -distribution,”⁴⁹ may possibly be regarded as a member of a broader kappa family of distributions, it seems incorrect to call it “the kappa distribution,” which is a well-established and much-used velocity/energy distribution function. In this context, we see that Bryant⁶ showed a particularly good example of a data fit, for solar wind electrons, to a conventional kappa distribution. The distribution clearly has more particles in both the low-energy and high-energy regions than a Maxwellian does, with a dip in the vicinity of the effective thermal speed. This would not fit the modified distributions used by Leubner^{45–47} and by Hau *et al.*,⁵⁰ which have fewer low-energy particles in a low- κ distribution than is the case for the Maxwellian.

Turning now to the paper by Hau and Fu,¹ we note that the distribution function is defined as

$$f^\kappa(v) = \frac{N}{2\pi(\kappa v_\kappa^2)^{3/2}} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa-1/2)\Gamma(\frac{3}{2})} \left(1 + \frac{v^2}{\kappa v_\kappa^2}\right)^{-(\kappa+1)}. \quad (13)$$

At first sight this is similar to our Eq. (9). However, although the characteristic speed v_κ appears from the notation (subscript κ) to have kappa dependence as θ does, it is, in fact, defined as $v_\kappa=[(2\kappa-3)/\kappa \times K_B T_\kappa/m]^{1/2}$, where “the characteristic temperature T_κ is $T_\kappa=\kappa T/(\kappa-3/2)$ with T being the temperature in the Maxwellian distribution, $f=N(m/2\pi K_B T)^{3/2} \exp(-mv^2/2K_B T)$.” By substituting for T_κ and hence for T in v_κ , one sees that this definition of the characteristic speed is circular, and that the denominator in Eq. (13) then simply reduces to $2\kappa K_B T/m = \kappa v_t^2$, i.e., $v_\kappa \equiv v_t$, the Maxwellian thermal speed. Thus, in this formulation, there is no need to introduce an apparently kappa-dependent characteristic speed v_κ .

The introduction of the notation T_κ in Ref. 1 appears to follow the notation of Maksimovic *et al.*,⁵¹ a paper cited in the Response to our Comment. However, the latter paper does not define T_κ in the form given in Ref. 1. Instead, it is given as $T_\kappa=m\langle v^2 \rangle/3K_B$, as in Eq. (8), i.e., they⁵¹ use the same form for the temperature as we have discussed above.^{3,14} As we have seen, that agrees with the form used in Ref. 10 and the many papers that follow it.

In the Response⁵⁰ to our Comment, the authors to some extent clarify their intention regarding v_κ , by explicitly writing the distribution in terms of the thermal velocity of a Maxwellian, rather than in terms of v_κ ,

$$f^\kappa(v) = \frac{N}{2\pi(\kappa v_{\text{th}}^2)^{3/2}} \frac{\Gamma(\kappa+1)}{\Gamma(\kappa-1/2)\Gamma(\frac{3}{2})} \left(1 + \frac{v^2}{\kappa v_{\text{th}}^2}\right)^{-(\kappa+1)}, \quad (14)$$

with v_{th} defined through $v_{\text{th}}^2=2K_B T/m$.

In principle, it does seem to be surprising, and indeed illogical, that a non-Maxwellian distribution should be defined *a priori* in terms of an unknown Maxwellian thermal speed. The result is that, in considering different values of κ , the authors are comparing different “equivalent temperatures” associated with different values of kappa with the temperature of a specific preordained Maxwellian. It is not clear

how that temperature is defined and to be measured. Bearing in mind the uncertainty of the meaning of a temperature for a non-Maxwellian (“nonequilibrium”) distribution, that would seem to be an exercise fraught with difficulties in interpretation. We believe that the standard approach, as we have outlined above, is both more satisfying theoretically and relates to the observable quantity, the mean kinetic energy per particle.

Furthermore, in using the thermal speed rather than the most probable speed θ , the authors¹ may appear to be following the model of Leubner.^{45–47} However, as we have spelt out above, the latter papers use a distribution in which the inverse power is κ , while in the authors’ representation¹ it is $(\kappa+1)$. Thus, the form of distribution function used by Hau and Fu is different from both the conventional form [Eq. (9)] and the Tsallis-like form [Eq. (12)].

Following on from their distribution [Eq. (13)], the authors obtain an expansion¹ of a normalized form of $f^\kappa(v)$ in the form of a power series in (x/κ) , where $x=(v/v_\kappa)^2$, as given in their Eqs. (3) and (4). However, as we have seen, x itself includes κ -dependence if v_κ is correctly defined, as given for θ above. Thus the expansion given by Hau and Fu¹ neglects dependence on κ that a proper definition of x would provide. One can see immediately from the formulation given in our Eq. (10) above that to rectify this it is important to write the expansion not in (x/κ) , but in $(X/[\kappa-3/2])$, where $X=v^2/(2K_B T/m)=(v/v_\kappa)^2$ is a variable that is independent of κ . We note that $X=x(\kappa-3/2)/\kappa$. This obviously has mathematical (and hence physical) implications down the line, such as the range of v over which the expansion is valid, for distributions with a very high superthermal content, i.e., for, say, $5/2 > \kappa > 3/2$.

Later in the paper,¹ the authors calculate moments of their distribution function to find expressions for the density and hence pressure of a species designated as α , in an electrostatic potential φ , to obtain

$$n_\alpha = N \left(1 + \frac{q_\alpha \varphi}{\kappa K_B T_\alpha} \right)^{-\kappa+1/2}, \quad (15)$$

in their notation. We stress that, as the denominator does not represent the full dependence of θ on κ , Eq. (15) suppresses some of the κ -dependence of the density in a potential φ . This is particularly important for low values of κ , approaching $(3/2)$. This construction is carried through from Eqs. (12) to (17) of their paper.¹

In fact, the full dependence of the density function on κ is reflected in our recent papers,^{32,33} where we have written (in equivalent notation) the density for electrons and ions in an electrostatic potential φ explicitly as

$$n_\alpha(\varphi) = N_\alpha \left(1 \pm \frac{2e\varphi}{m_\alpha \kappa_\alpha \theta_\alpha^2} \right)^{-(\kappa_\alpha-1/2)}, \quad (16)$$

with $\theta_\alpha^2 = [(\kappa_\alpha-3/2)/\kappa_\alpha][2K_B T_\alpha/m_\alpha]$ and $\alpha=e, i$ for the respective species. In this expression the upper (lower) sign gives the expression for the ions (electrons) and T_α is the temperature of the equivalent Maxwellian, as discussed above. Substitution for θ_α then reveals the full dependence of the density on κ_α , viz.,

$$n_\alpha(\varphi) = N_\alpha \left(1 \pm \frac{e\varphi}{[\kappa_\alpha-3/2]K_B T_\alpha} \right)^{-(\kappa_\alpha-1/2)}. \quad (17)$$

Naturally, this expression is obtained from either the three-dimensional or the one-dimensional form of the kappa distribution function.³² Appropriate normalized forms may then be deduced for each species,³² characterized by the factor $(\kappa_\alpha-3/2)$ in the denominator.

Comparing Eq. (17) with that obtained by Hau and Fu^{1,50} [Eq. (15)], one sees that the latter does not reflect the rapid changes that occur for κ values approaching $(3/2)$.

Finally, we note that in their Response⁵⁰ the authors cite a paper by Collier.⁵² However, a careful reading of that work shows that Collier used the form of kappa distribution given in Ref. 10, i.e., our Eq. (9), and then indicated that one normally defines a temperature for this non-Maxwellian distribution in the form that we have given above [Eq. (8)]. Collier then argued that in evaluating such a temperature from data, there are at times difficulties in fitting the low-flux high-energy part of the spectrum, resulting in the fit often being dominated by the core Maxwellian part. On that basis he introduced a second temperature T_{core} , associated with that Maxwellian core. Interestingly, he found that for ion distributions T_{core} increases with increasing κ , a result which is in line with our earlier discussion concerning the narrowness of the “core” distribution of a kappa distribution.

In related papers,^{53,54} the kappa distribution is again presented in the conventional form, Eq. (9), although the core temperature is also introduced in the discussion. We also note that a close examination of Fig. 2 of Ref. 55 indicates that the calculated κ distribution found by Collier in that case appears to have a central peak that is higher than that of the Maxwellian, i.e., it agrees better with the shape obtained from Eq. (9) than with that from the alternative distribution, as given in Fig. 1 of the Response.⁵⁰

In summary, we agree that Hau and Fu¹ have found a number of useful mathematical expressions for a non-Maxwellian distribution that, like a kappa distribution, has a high-energy tail. However, we have shown that in their calculations they have used neither the standard, conventional kappa distribution [Eq. (9)] (Refs. 2, 10, and 13) nor the modified (q -like) distribution [Eq. (12)] recently suggested,⁴⁵ that is based on nonextensive entropy. Further we have shown that as a result of an unusual definition of T_κ , aspects of their original presentation¹ involve a circular argument, as a result of which their introduction of v_κ is superfluous. We have drawn attention to the misleading nature of the expansion that they obtain for low velocities, in that it suppresses an important part of the dependence of the kappa distribution function on κ , and we have given the density of a κ -distributed component in an electrostatic potential, in a form in which its full dependence on κ is reflected.

This research is supported in part by the National Research Foundation of South Africa (NRF). Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and therefore the NRF does not accept any liability in regard thereto. T.K.B. acknowledges with thanks partial funding from DST through NASSP.

N.S.S. would like to thank Guru Nanak Dev University, Amritsar, India for providing leave. The work of N.S.S. and I.K. was supported by EPSRC-UK Science and Innovation award to the Centre for Plasma Physics (QUB), CPP Grant No. EP/D06337X/1.

- ¹L.-N. Hau and W.-Z. Fu, *Phys. Plasmas* **14**, 110702 (2007).
- ²V. M. Vasyliunas, *J. Geophys. Res.* **73**, 2839, doi:10.1029/JA073i009p02839 (1968).
- ³V. Formisano, G. Moreno, F. Palmiotto, and P. C. Hedgecock, *J. Geophys. Res.* **78**, 3714, doi:10.1029/JA078i019p03714 (1973).
- ⁴S. P. Christon, D. G. Mitchell, D. J. Williams, L. A. Frank, C. Y. Huang, and T. E. Eastman, *J. Geophys. Res.* **93**, 2562, doi:10.1029/JA093iA04p02562 (1988).
- ⁵W. C. Feldman, R. C. Anderson, J. R. Asbridge, S. J. Bame, J. T. Gosling, and R. D. Zwickl, *J. Geophys. Res.* **87**, 632, doi:10.1029/JA087iA02p00632 (1982); W. C. Feldman, R. C. Anderson, S. J. Bame, S. P. Gary, J. T. Gosling, D. J. McComas, M. F. Thomsen, G. Paschmann, and M. M. Hoppe, *ibid.* **88**, 96, doi:10.1029/JA088iA01p00096 (1983).
- ⁶D. A. Bryant, *J. Plasma Phys.* **56**, 87 (1996).
- ⁷A. F. Viñas, R. L. Mace, and R. F. Benson, *J. Geophys. Res.* **110**, A06202, doi:10.1029/2004JA010967 (2005).
- ⁸P. Schippers, M. Blanc, N. André, I. Dandouras, G. R. Lewis, L. K. Gilbert, A. M. Persoon, N. Krupp, D. A. Gurnett, A. J. Coates, S. M. Krimigis, D. T. Young, and M. K. Dougherty, *J. Geophys. Res.* **113**, A07208, doi:10.1029/2008JA013098 (2008).
- ⁹M. A. Hellberg, R. L. Mace, R. J. Armstrong, and G. Karlstad, *J. Plasma Phys.* **64**, 433 (2000).
- ¹⁰D. Summers and R. M. Thorne, *Phys. Fluids B* **3**, 1835 (1991).
- ¹¹D. Summers and R. M. Thorne, *J. Geophys. Res.* **95**, 1133, doi:10.1029/JA095iA02p01133 (1990).
- ¹²N. A. Krall and A. W. Trivelpiece, *Principles of Plasma Physics* (McGraw-Hill, New York, 1973), p. 6.
- ¹³M. P. Leubner, *J. Geophys. Res.* **87**, 6335, doi:10.1029/JA087iA08p06335 (1982).
- ¹⁴Y. F. Chateau and N. Meyer-Vernet, *J. Geophys. Res.* **96**, 5825, doi:10.1029/90JA02565 (1991).
- ¹⁵R. M. Thorne and D. Summers, *Phys. Fluids B* **3**, 2117 (1991).
- ¹⁶R. M. Thorne and D. Summers, *J. Geophys. Res.* **96**, 217, doi:10.1029/90JA01629 (1991).
- ¹⁷Z. Meng, R. M. Thorne, and D. Summers, *J. Plasma Phys.* **47**, 445 (1992).
- ¹⁸S. Xue, R. M. Thorne, and D. Summers, *J. Geophys. Res.* **98**, 17475, doi:10.1029/93JA00790 (1993).
- ¹⁹D. Summers, S. Xue, and R. M. Thorne, *Phys. Plasmas* **1**, 2012 (1994).
- ²⁰B. D. Fried and S. D. Conte, *The Plasma Dispersion Function* (Academic, New York, 1961).
- ²¹R. L. Mace and M. A. Hellberg, *Phys. Plasmas* **2**, 2098 (1995).
- ²²R. L. Mace, *J. Plasma Phys.* **55**, 415 (1996).
- ²³C. Grabbe and E. Venturino, *Phys. Plasmas* **3**, 35 (1996).
- ²⁴R. L. Mace, *J. Geophys. Res.* **103**, 14643, doi:10.1029/98JA00616 (1998).
- ²⁵R. L. Mace, M. A. Hellberg, and R. A. Treumann, *J. Plasma Phys.* **59**, 393 (1998).
- ²⁶R. L. Mace, G. Amery, and M. A. Hellberg, *Phys. Plasmas* **6**, 44 (1999).
- ²⁷M. A. Hellberg and R. L. Mace, *Phys. Plasmas* **9**, 1495 (2002).
- ²⁸R. L. Mace and M. A. Hellberg, *Phys. Plasmas* **10**, 21 (2003).
- ²⁹R. L. Mace, *Phys. Plasmas* **10**, 2181 (2003).
- ³⁰R. L. Mace, *Phys. Plasmas* **11**, 507 (2004).
- ³¹T. Cattaert, M. A. Hellberg, and R. L. Mace, *Phys. Plasmas* **14**, 082111 (2007).
- ³²T. K. Baluku and M. A. Hellberg, *Phys. Plasmas* **15**, 123705 (2008).
- ³³N. S. Saini, I. Kourakis, and M. A. Hellberg, *Phys. Plasmas* **16**, 062903 (2009).
- ³⁴Y.-D. Jung, *Phys. Lett. A* **313**, 296 (2003).
- ³⁵C.-G. Kim and Y.-D. Jung, *Plasma Phys. Controlled Fusion* **46**, 1493 (2004).
- ³⁶C.-G. Kim and Y.-D. Jung, *J. Phys. B* **38**, 3867 (2005).
- ³⁷J. J. Podesta, "Plasma dispersion function for the kappa distribution," NASA Report No. NASA/CR-2004-212770, 2004.
- ³⁸J. J. Podesta, *Phys. Plasmas* **12**, 052101 (2005).
- ³⁹M. A. Raadu and M. Shafiq, *Phys. Plasmas* **14**, 012105 (2007).
- ⁴⁰M. A. Sarwar, A. M. Mirza, and M. S. Qaisara, *Phys. Plasmas* **14**, 073702 (2007).
- ⁴¹M.-J. Lee, *Phys. Plasmas* **14**, 032112 (2007).
- ⁴²M. Lazar, R. Schlickeiser, and P. K. Shukla, *Phys. Plasmas* **15**, 042103 (2008).
- ⁴³M. Lazar, R. Schlickeiser, S. Poedts, and R. C. Tautz, *Mon. Not. R. Astron. Soc.* **390**, 168 (2008).
- ⁴⁴B. Basu, *Phys. Plasmas* **15**, 042108 (2008).
- ⁴⁵M. P. Leubner, *Astrophys. Space Sci.* **282**, 573 (2002).
- ⁴⁶M. P. Leubner, *Phys. Plasmas* **11**, 1308 (2004).
- ⁴⁷M. P. Leubner, *Astrophys. J.* **604**, 469 (2004).
- ⁴⁸R. Silva, Jr., A. R. Plastino, and J. A. S. Lima, *Phys. Lett. A* **249**, 401 (1998).
- ⁴⁹C. Tsallis, *Physica A* **221**, 277 (1995).
- ⁵⁰L.-N. Hau, W.-Z. Fu, and S.-H. Chuang, *Phys. Plasmas* **16**, 094702 (2009).
- ⁵¹M. Maksimovic, V. Pierrard, and P. Riley, *Geophys. Res. Lett.* **24**, 1151, doi:10.1029/97GL00992 (1997).
- ⁵²M. R. Collier, *Geophys. Res. Lett.* **22**, 2673, doi:10.1029/95GL02350 (1995).
- ⁵³M. R. Collier and D. C. Hamilton, *Geophys. Res. Lett.* **22**, 303, doi:10.1029/94GL02997 (1995).
- ⁵⁴M. R. Collier, *J. Geophys. Res.* **104**, 28559, doi:10.1029/1999JA900355 (1999).
- ⁵⁵M. R. Collier, *Adv. Space Res.* **33**, 2108 (2004).